


ANNUAL SUMMARY 1978
(REVISED)
REISSUED FEBRUARY 1981

CENTERS FOR DISEASE CONTROL
FOODBORNE DISEASE

SURVEILLANCE



PREFACE

This report summarizes information received from state and local health departments, the Food and Drug Administration, the U.S. Department of Agriculture, and other pertinent sources. The information is preliminary and is intended primarily for use by those with responsibility for disease control activities. Anyone desiring to quote this report should contact the Enteric Diseases Branch for confirmation and further interpretation.

Contributions to the report are most welcome. Please address them to:

Centers for Disease Control
ATTN: Bureau of Epidemiology
Bacterial Diseases Division
Enteric Diseases Branch
Atlanta, Georgia 30333

SUGGESTED CITATION

Centers for Disease Control: Foodborne Disease Outbreaks
Annual Summary 1978, Revised
Reissued February 1981

- Centers for Disease Control William H. Foege, M.D., Director
- Center for Infectious Diseases Walter R. Dowdle, Ph.D., Director
- Bacterial Diseases Division John V. Bennett, M.D., Director
- Enteric Diseases Branch Roger A. Feldman, M.D., Chief
- Foodborne Disease Outbreaks
Surveillance Activity John D. Snyder, M.D.
- Statistical Services Branch Stanley M. Martin, M.S., Chief
Robert A. Pollard, M.A.
Dwan G. Smith
- Epidemiologic Investigations
- Laboratory Branch George K. Morris, Ph.D., Chief
- Enteric Diseases Laboratory Section Joy G. Wells, M.S., Chief
- Editorial and Graphic Services Frances H. Porcher, M.A., Chief
Charlotte C. Turner, Writer-Editor
- Bureau of Laboratories Roslyn Q. Robinson, Ph.D., Director
- Bacteriology Division Albert Balows, Ph.D., Director
- Enterobacteriology Branch V. R. Dowell, Jr., Ph.D., Chief
- Clinical Bacteriology Branch P. B. Smith, Ph.D., Chief
- Bacterial Immunology Branch John C. Feeley, Ph.D., Chief

* This report has been reissued because of numerous tabular errors in the original 1978 *
* Foodborne Disease Surveillance Report released in November 1979. The data from the *
* previous 1978 report should be disregarded and copies should be discarded. *

TABLE OF CONTENTS

	Page
I. INTRODUCTION	1
II. FOODBORNE DISEASE OUTBREAKS	1
A. Definition of Outbreak	2
B. Source of Data	2
C. Interpretation of Data	2
D. Analysis of Data	4
E. Investigation of a Foodborne Outbreak (Sample Form)	17
F. Line Listing of Foodborne Disease Outbreaks, 1978	19
G. Guidelines for Confirmation of Foodborne Outbreak	42
H. Selected Foodborne Outbreak Articles, 1978, Taken From Morbidity and Mortality Weekly Report	47
I. Bibliography	50



I. INTRODUCTION

The reporting of foodborne and waterborne diseases in the United States began over half a century ago when state and territorial health officers, concerned about the high morbidity and mortality caused by typhoid fever and infantile diarrhea, recommended that cases of enteric fever be investigated and reported. The purpose was to obtain information about the role of food, milk, and water in outbreaks of intestinal illness as the basis for sound public health action. Beginning in 1923, the United States Public Health Service published summaries of outbreaks of gastrointestinal illness attributed to milk. In 1938, it added summaries of outbreaks caused by all foods. These early surveillance efforts led to the enactment of important public health measures which had a profound influence in decreasing the incidence of enteric diseases, particularly those transmitted by milk and water.

From 1951 through 1960, the National Office of Vital Statistics reviewed reports of outbreaks of foodborne illness and published summaries of them annually in Public Health Reports. In 1961 the Center for Disease Control (CDC), then the Communicable Disease Center, assumed responsibility for publishing reports on foodborne illness. For the period 1961-66 CDC discontinued publication of annual reviews, but reported pertinent statistics and detailed individual investigations in the Morbidity and Mortality Weekly Report (MMWR).

In 1966 the present system of surveillance of foodborne and waterborne diseases began with the incorporation of all reports of enteric disease outbreaks attributed to microbial or chemical contamination of food or water into an annual summary. Since 1966 the quality of investigative reports has improved primarily as a result of more active participation by state and federal agencies in the investigation of foodborne and waterborne disease outbreaks. Due to increasing interest and activity in waterborne disease surveillance, foodborne and waterborne disease outbreaks were reported in separate annual summaries for the first time in 1978. This report summarizes data from the foodborne disease outbreaks reported to the CDC in 1978.

Foodborne disease surveillance has traditionally served 3 objectives:

1. Disease Prevention and Control: Early identification and removal of contaminated products from the commercial market, correction of faulty food preparation practices in food service establishments and in the home, and identification and appropriate treatment of human carriers of foodborne pathogens are the fundamental prevention and control measures resulting from surveillance of foodborne disease.

2. Knowledge of Disease Causation: The responsible pathogen was not identified in 30% to 60% of foodborne disease outbreaks reported to CDC in each of the last 5 years. In many of these outbreaks, pathogens known to cause foodborne illness may not have been identified because of late or incomplete laboratory investigation. In others, the responsible pathogen may have escaped detection even when a thorough laboratory investigation was carried out because the pathogen is not yet appreciated as a cause of foodborne disease or because it cannot yet be identified by available laboratory techniques. It is probable that these pathogens can be identified and suitable measures to prevent or control diseases caused by them can be instituted if more thorough clinical, epidemiologic, and laboratory investigations are employed. Pathogens suspected of being, but not yet determined to be, etiologic agents in foodborne disease include Group D Streptococcus, Citrobacter, Enterobacter, Klebsiella, Pseudomonas, and the presumably viral agents of acute infectious non-bacterial gastroenteritis. Other pathogens such as Escherichia coli, Bacillus cereus, Yersinia enterocolitica, Vibrio parahaemolyticus and Campylobacter fetus subspecies jejuni are known causes of foodborne illness, but the extent and importance of their role have not been adequately assessed as yet.

3. Administrative Guidance: The collection of data from outbreak investigations permits assessment of trends in etiologic agents and food vehicles and focuses on common errors in food handling. By compiling the data in an annual summary, it is hoped that local and state health departments and others involved in the implementation of food protection programs will be kept informed of the factors involved in foodborne disease outbreaks. Comprehensive surveillance would result in a clearer appreciation of priorities in food protection, institution of better training programs, and more rational utilization of available resources.

II. FOODBORNE DISEASE OUTBREAKS

In 1978, 481 outbreaks of foodborne disease involving 10,639 cases were reported to the Centers for Disease Control.

A. Definition of Outbreak

For the purpose of this report, a foodborne disease outbreak is defined as an incident in which (1) 2 or more persons experience a similar illness, usually gastrointestinal, after ingestion of a common food, and (2) epidemiologic analysis implicates the food as the source of the illness. There are a few exceptions; 1 case of botulism or chemical poisoning constitutes an outbreak.

1. Laboratory confirmed--Outbreaks in which laboratory evidence of a specific etiologic agent is obtained, and specified criteria are met (see Section G).

2. Undetermined etiology--Outbreaks in which epidemiologic evidence implicates a food source, but adequate laboratory confirmation is not obtained. These outbreaks are subdivided into 4 subgroups by incubation period of the illnesses: less than 1 hour (probable chemical poisoning), 1 to 7 hours (probable Staphylococcus food poisoning), 8 to 14 hours (probable Clostridium perfringens), and greater than 14 hours (infectious or toxic agents).

B. Source of Data

The general public and local, state, and federal agencies which have responsibility for public health and food protection participate in foodborne disease surveillance. Consumers, physicians, hospital personnel, and persons involved with food service or processing report complaints of illness to health departments or regulatory agencies. Local health department personnel (including epidemiologists, sanitarians, and public health nurses) carry out most epidemiologic investigations of these reports and make their findings available to state health departments. State agencies concerned with food safety frequently participate in the initial investigation of the outbreak and offer laboratory support. State or other officials eventually summarize the findings of the investigation on the standard CDC reporting form (see Section E) and send these to CDC (Table 1). Occasionally, on special request, CDC participates in an investigation, particularly if the outbreak is large or involves products that move in interstate commerce.

The 2 federal regulatory agencies that have major responsibilities for food protection, the Food and Drug Administration (FDA) and U.S. Department of Agriculture (USDA), report episodes of foodborne illness to CDC and to state and local health authorities which, in turn, report to FDA or USDA any foodborne disease outbreaks involving commercial products. The U.S. Armed Forces also report outbreaks directly to CDC.

By special arrangement, Connaught Laboratories of Canada, the only commercial producer of botulinum antitoxin in the Western Hemisphere, immediately report all requests for botulinum antitoxin to CDC. This is sometimes the first communication of a botulism outbreak to public health authorities, although physicians are urged to promptly report all suspected botulism cases. In botulism outbreaks, CDC works closely with physicians, state and local health authorities, and FDA or USDA representatives to provide diagnostic and therapeutic consultation and to rapidly identify the responsible food or foods so that proper corrective action can be taken.

C. Interpretation of Data

The limitations on the quantity and quality of data presented here must be appreciated in order to avoid misinterpretation. The number of outbreaks of foodborne disease reported by this surveillance system clearly represents a small fraction of the total number that occur. The likelihood of an outbreak coming to the attention of health authorities varies considerably depending on consumer and physician awareness, interest, and motivation to report the incident.

Not all cases of foodborne illness have an equal likelihood of being reported. For example, interstate outbreaks, large intrastate outbreaks, and outbreaks of serious illness such as botulism or amanitotoxin (mushroom) poisoning are more likely to come to the attention of health authorities than cases of mild illness following a family cookout.

The quality of the data presented here depends upon the commitment given to foodborne surveillance by the state or local health departments. Not only the department's interest in foodborne disease investigation but its investigative and laboratory capabilities are essential determinants of the quality of the investigation. Similarly, the likelihood that the findings of an investigation will be reported varies from one locale to another.

Just as this report should not be the basis of firm conclusions about the absolute incidence of foodborne disease, it should not be used to draw conclusions about the relative incidence of foodborne disease of various etiologies (Table 2). For example, foodborne diseases characterized by short incubation periods such as most outbreaks of chemical etiology or outbreaks caused by staphylococcal enterotoxin are more likely to be recognized

as common-source foodborne disease outbreaks than those diseases with longer incubation periods such as hepatitis A. The relatively small number of outbreaks attributed to parasites and viruses may be due to the long incubation periods of these agents masking the common-source nature of many cases. Similarly, outbreaks involving B. cereus, E. coli, V. parahaemolyticus, Y. enterocolitica, or C. fetus ssp jejuni are probably less likely to be confirmed because these organisms are often not considered in clinical, epidemiologic, and laboratory investigations.

The number of reported outbreaks attributed to some etiologies depends upon the interest of a particular health department or individual. For example, the cholera cases uncovered in Louisiana in 1978 might have been missed if it had not been for the persistence of a laboratory technician who sought aid in identifying an organism which he could not type.

Establishing the true number of deaths caused by foodborne disease outbreaks is difficult because information on deaths in the reports is often incomplete or absent. Further contributing to the under-reporting is the fact that foodborne disease may not be recognized as contributing to the demise of an elderly or debilitated person unable to withstand otherwise minor physical stresses. These limitations on the data must be understood in interpreting Table 3.

In outbreaks of unknown etiology, listed by incubation period (Table 4), the accuracy of reported information is always suspect. In these outbreaks when the epidemiology incriminating a particular food item was very weak, the food was listed as unknown in this report (Table 5). Previously, persons with botulism for which no vehicle could be implicated were included in the foodborne totals. However, in 1978 the definition of foodborne botulism was restricted to cases where a food source was confirmed either by laboratory or epidemiologic evidence. In addition to the 12 foodborne botulism outbreaks, there were 10 botulism outbreaks in 1978 involving 13 persons where no food or wound source could be found.

Information on the place of eating suspect food in foodborne outbreaks was judged to be reliable and was recorded (Table 6). However, information on the place where food was mishandled or improperly cooked or stored in these outbreaks was generally judged unreliable; in many of them the place of mishandling was listed as unknown (Table 7). Only in outbreaks in which a specific etiology was highly suspected, although unconfirmed in the laboratory, and in which the information on mishandling was consistent with the suspected etiology was a known place of mishandling designated.

The implications of a food-processing establishment mishandling food are great both to public health authorities and the establishment concerned. Consequently, the outbreaks attributed to mishandling at these establishments are thoroughly investigated and reported data carefully scrutinized. No foodborne disease outbreaks were linked to mishandling of food at a food processing plant in 1978.

Much is known about contributing factors in foodborne disease. The five most common factors contributing to foodborne disease outbreaks in the United States in order of frequency of occurrence include (1) inadequate cooling of foods, (2) lapse of a day or more between preparing and serving, (3) infected persons handling foods which are not subsequently heat-processed, (4) inadequate time or temperature or both during heat processing foods, and (5) insufficiently high temperature during storage of hot foods (1). For example, in most outbreaks of botulism and trichinosis, the food is usually inadequately heat treated. In most of the outbreaks of bacterial etiology other than botulism and in outbreaks of scrombroid (in which bacterial growth is responsible for toxin production), the food is usually stored at improper holding temperatures. By definition, in outbreaks of ciguatera, puffer fish poisoning, mushroom poisoning, and paralytic and neurotoxic shellfish poisoning, the food itself is unsafe, and illness is not related to improper handling or preparation.

The investigators of foodborne disease outbreaks are usually aware of these contributing factors and consequently seek and find the appropriate answers. Sometimes, however, investigators report factors which are not known to be contributing to outbreaks of the type of etiology confirmed. In such cases the factors are considered in light of the evidence presented; if they are totally unsubstantiated, they are rejected. These considerations must be borne in the mind in interpreting Table 8.

Reference

1. Bryan FA. Factors that contribute to outbreaks of foodborne disease. J Food Protection 1978;41:816-827.

D. Analysis of Data

In 1978, 481 outbreaks of foodborne disease involving 10,639 cases were reported to the CDC Foodborne Disease Surveillance Activity, compared with the 5-year average of 426 outbreaks and 13,709 outbreaks from 1973 to 1977. Outbreaks were reported from 42 states, New York City, Puerto Rico, Guam, and the Virgin Islands. No outbreaks were reported from 8 states or the District of Columbia.

The large number of outbreaks reported by several states and New York City undoubtedly reflects the interest and effort at control of the respective health departments in foodborne disease surveillance. The Washington State Department of Social and Health Services has continued its record of reporting more outbreaks than any other state (Table 1). California, Pennsylvania, and Hawaii again are among the leading states in reporting outbreaks of foodborne disease. New York City has reported more foodborne outbreaks than any other reporting agency since 1975. Connecticut, Maryland, New Mexico, and Virginia reported a substantial increase in the number of outbreaks in 1978.

As has been seen in each of the 5 preceeding years, bacterial agents were the most common causes (68%) of the foodborne outbreaks of confirmed etiology. Following in frequency were chemical agents (24%), parasitic (5%), and viral (3%). The overwhelming majority of confirmed cases (90%) were of bacterial etiology, nearly matching the 5-year average of 91%. *Salmonellae* accounted for about one-third of the confirmed outbreaks and nearly 40% of the cases, figures consistent with results for 1973 to 1977. The second most common agent was *Staphylococcus aureus* which was implicated in 15% of the outbreaks and 27% of the total cases.

For the first year since the beginning of foodborne disease reporting, an outbreak of *Vibrio cholerae* serotype O1 disease was detected in the United States (1). In addition to 1 cluster involving 4 persons, 7 additional individual cases were uncovered, all in Louisiana, with boiled crabs identified as the vehicle.

In 1978, 14 deaths were reported as associated with foodborne outbreaks (Table 3). Four deaths were caused by hepatitis, 3 by botulism, and 7 were caused by diseases of unknown etiology.

The etiologic agent was confirmed in 32% (154/481) of the outbreaks, which is slightly lower than the 5-year average of 40%. Table 4 lists the outbreaks of diseases of undetermined etiology by median incubation periods. If one assumes that most outbreaks in which the median incubation period was less than 1 hour were chemical poisoning, that those in which median incubation period was 1-7 hours were of staphylococcal intoxication, and that those in which the median incubation period was 8-14 hours were caused by *C. perfringens*, then 60% of the outbreaks of unknown etiology can be accounted for. In addition to these agents, *B. cereus*, which is rarely considered either from an epidemiologic standpoint or in the laboratory, is associated with 2 separate food poisoning syndromes which closely resemble the more familiar ones caused by *S. aureus* and *C. perfringens* (2). Determination of the relative importance of *B. cereus* infections must await an increased awareness of the potential of the organism to cause foodborne illness.

The vehicle of transmission was identified in 81% of the 154 outbreaks of known etiology. The most common vehicles were meat (39/154) and fish and shellfish (34/154) in outbreaks of bacterial etiology for which a single food item could be identified.

Home canned foods were the most frequently incriminated vehicle in outbreaks of botulism. However, potato salad prepared in commercial food establishments was implicated in 2 outbreaks involving 45 of the 58 cases. *Salmonella* outbreaks were caused by a variety of vehicles including meat, poultry, dairy products, salads, and Mexican food. Ciguatoxin, the third most frequently reported confirmed agent, caused outbreaks involving mainly coral reef fish. All the outbreaks of paralytic shellfish poisoning were associated with scallops or mussels.

Food eaten in the home (122/481) and restaurants (234/481) accounted for 74% of the 1978 foodborne outbreaks. Of the 105 bacterial outbreaks, 39 were attributed to food eaten in the home and 32 in restaurants. Chemical outbreaks were more likely to occur in the home (21 of 37), and 6 of the parasitic outbreaks occurred in the home.

In 1978 no reports of foodborne outbreaks due to mishandling or improper cooking and storage of food at food processing establishments were reported (Table 7). Mishandling of food at food service establishments accounted for 28% of the outbreaks while mishandling at home was implicated in 8%.

Errors in food handling practices responsible for outbreaks were reported in 75 of the 154 outbreaks of known etiology (Table 8). Improper holding temperatures or inadequate cooking were responsible for most of the outbreaks of bacterial etiology. Poor personal hygiene of a food handler was also frequently reported as a contributing factor, especially

in foodborne shigellosis and in viral hepatitis outbreaks. Inadequate cooking was a factor in all outbreaks due to parasites in which contributing factors were reported.

Since the toxins responsible for ciguatera, mushroom, and paralytic shellfish poisoning are heat stable, thorough cooking of food does not provide protection from these illnesses. Furthermore, there is no practical way to distinguish fish or shellfish containing ciguatoxin or neurotoxin. For these reasons, a place of food mishandling was not specified in outbreaks of ciguatera, mushroom, or shellfish poisoning.

In reviewing the 481 outbreaks, at least 1 contributing factor was implicated in 206 (43%) (Table 8). The data mirrored patterns seen the previous 5 years. In reported outbreaks of botulism, the most frequent error was inadequate cooking of food. Improper holding temperatures most frequently contributed to reported outbreaks of salmonellosis, staphylococcus intoxication, and C. perfringens foodborne illness. Heavy metal poisoning was usually due to storage of acidic beverages in containers or pipes from which metal ions could be leached. In outbreaks of ciguatera, paralytic shellfish, and mushroom poisoning, the foods were unsafe to begin with because they contained toxins.

The date of onset of an outbreak was designated as the date of onset of the first case (Table 9). Certain types of foodborne disease outbreaks showed a definite seasonality (Table 9). For example, outbreaks of paralytic shellfish poisoning occurred in September and October following the peak period of growth of dinoflagellates in the warm summer months.

References

1. Blake PA, Allegra DT, Snyder JD, Barrett TJ, et al. Cholera--a possible endemic focus in the United States. *N Engl J Med* 1980;302:305-9.
2. Terranova W, Blake PA. Bacillus cereus food poisoning. *N Engl J Med* 1978; 298:143-4.

Fig. 1 REPORTED FOODBORNE DISEASE OUTBREAKS, 1978




Table 1
Foodborne Disease Outbreaks, United States, by Location, 1978

<u>State</u>	<u>Number of Outbreaks</u>	<u>State</u>	<u>Number of Outbreaks</u>
Alabama	1	New Jersey	10
Alaska	1	New Mexico	24
Arizona	1	New York City	118
Arkansas	4	New York State	4
California	38	North Carolina	4
Colorado	6	North Dakota	0
Connecticut	24	Ohio	0
Delaware	0	Oklahoma	3
D.C.	0	Oregon	7
Florida	13	Pennsylvania	26
Georgia	14	Puerto Rico	2
Hawaii	22	Rhode Island	2
Idaho	4	South Carolina	2
Illinois	4	South Dakota	1
Indiana	1	Tennessee	1
Iowa	0	Texas	4
Kansas	6	Utah	2
Kentucky	1	Vermont	1
Louisiana	3	Virginia	8
Maine	3	Washington	51
Maryland	18	West Virginia	0
Massachusetts	0	Wisconsin	8
Michigan	2	Wyoming	0
Minnesota	13	Virgin Islands	3
Mississippi	0	Guam	4
Missouri	5	Total	481
Montana	4		
Nebraska	5		
Nevada	1		
New Hampshire	2		

Table 2
Confirmed Foodborne Disease Outbreaks and Cases, and
Percents of Known Etiology, 1978

<u>Etiology</u>	<u>Number of Outbreaks</u>	<u>%</u>	<u>No. of Cases</u>	<u>%</u>
<u>BACTERIAL</u>				
<u>B. cereus</u>	6	3.9	248	5.0
<u>Brucella</u>	-	0.0	-	0.0
<u>C. botulinum</u>	12	7.8	58	1.2
<u>C. perfringens</u>	9	5.8	617	12.4
<u>E. coli</u>	1	0.6	35	0.7
<u>Salmonella</u>	45	29.2	1921	38.7
<u>Shigella</u>	4	2.6	159	3.2
<u>S. aureus</u>	23	14.9	1318	26.6
<u>Enterococci</u>	1	0.6	5	0.1
<u>Streptococcus Group A</u>	-	0.0	-	0.0
<u>V. cholerae 01</u>	1	0.6	11	0.2
<u>V. parahaemolyticus</u>	2	1.3	86	1.7
<u>Other bacterial</u>	1	0.6	8	0.2
Total	105	68.2	4466	90.0
<u>CHEMICAL</u>				
Heavy metals	1	0.6	41	0.8
Ciguatoxin	19	12.3	56	1.1
Neurotoxic shellfish poisoning	-	0.0	-	0.0
Paralytic shellfish poisoning	4	2.6	10	0.2
Scrombrotoxin	7	4.5	30	0.6
Monosodium glutamate	-	0.0	-	0.0
Mushroom poisoning	1	0.6	7	0.1
<u>Other chemicals</u>	5	3.2	19	0.4
Total	37	24.0	163	3.3
<u>PARASITIC</u>				
<u>T. spiralis</u>	7	4.5	35	0.7
Total	7	4.5	35	0.7
<u>VIRAL</u>				
Hepatitis A	5	3.2	300	6.0
Total	5	3.2	300	6.0
 CONFIRMED TOTAL	 154	 100.0	 4964	 100.0

Table 3

Deaths Associated with Foodborne Disease Outbreaks, 1978

<u>Etiology</u>	<u>Number of Deaths</u>
<u>C. botulinum</u>	3
Hepatitis A	4
Unknown	<u>7</u>
Total	14

Table 4

Foodborne Disease Outbreaks of Unknown
Etiology, by Incubation Period, 1978

<u>Incubation Period</u>	<u>Number of Outbreaks</u>
<1 hour	13
1-7 hours	109
8-14 hours	74
>15 hours	62
Unknown	<u>69</u>
Total	327

Table 5
Foodborne Disease Outbreaks, by Vehicle of Transmission, and Specific Etiology, 1978

<u>Etiology</u>	<u>Beef</u>	<u>Veal</u>	<u>Lamb</u>	<u>Ham</u>	<u>Pork</u>	<u>Sau- sage</u>	<u>Chicken</u>	<u>Turkey</u>	<u>Other Meat</u>	<u>Shell Fish</u>	<u>Crus- tacea</u>	<u>Other Fish</u>	<u>Eggs</u>	<u>Milk</u>
<u>BACTERIAL</u>														
<u>B. cereus</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Brucella</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>C. botulinum</u>	-	-	-	-	-	-	-	-	1	-	-	1	-	-
<u>C. perfringens</u>	2	-	-	-	1	-	-	-	-	-	-	-	-	-
<u>E. coli</u>	1	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Salmonella</u>	3	-	-	-	7	-	1	3	3	-	-	-	-	2
<u>Shigella</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>S. aureus</u>	-	-	-	10	-	-	-	-	-	-	-	-	-	-
<u>Enterococci</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Streptococcus Group A</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>V. cholerae</u>	-	-	-	-	-	-	-	-	-	1	-	-	-	-
<u>V. parahaemolyticus</u>	-	-	-	-	-	-	-	-	-	2	-	-	-	-
<u>Other bacterial</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>CHEMICAL</u>														
<u>Heavy metals</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Ciguatoxin</u>	-	-	-	-	-	-	-	-	-	-	-	19	-	-
<u>Neurotoxic shellfish</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Paralytic shellfish</u>	-	-	-	-	-	-	-	-	-	4	-	-	-	-
<u>Scrombrotoxin</u>	-	-	-	-	-	-	-	-	-	-	-	7	-	-
<u>Monosodium glutamate</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Mushroom poisoning</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>Other chemicals</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>PARASITIC</u>														
<u>T. spiralis</u>	3	-	-	-	1	1	-	-	2	-	-	-	-	-
<u>VIRAL</u>														
<u>Hepatitis A</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>CONFIRMED TOTAL</u>	9	-	-	10	9	1	1	3	6	7	-	27	-	2
<u>UNKNOWN</u>	5	-	-	2	1	1	3	-	2	9	-	2	-	-
<u>TOTAL</u>	14	-	-	12	10	2	4	3	8	16	-	29	-	2

Continued on page 11

Table 5 (Cont'd)
 Foodborne Disease Outbreaks, by Vehicle of Transmission, and Specific Etiology, 1978

Cheese	Egg Nog	Ice Cream	Non Dar Bev	Other Dairy	Baked Foods	Fruits &Vege- tables	Potato Salad	Egg Salad	Other Salads	Mush- rooms	Chi- nese Food	Mexi- can Food	Car- bonated Bev	Multi- ple Foods	Other Foods	Unknown	Total
-	-	-	-	-	-	-	-	-	-	-	2	1	-	-	1	2	6
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
-	-	-	-	-	-	4	1	-	1	-	-	1	-	1	2	-	12
-	-	-	-	-	-	-	-	-	1	-	-	2	-	1	-	2	9
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1
-	1	1	-	1	-	-	1	-	1	-	-	2	-	3	1	15	45
-	-	-	-	-	-	-	1	-	1	-	-	-	-	-	-	2	4
-	-	-	-	-	-	-	1	1	5	-	-	-	-	4	-	2	23
-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	1
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	1
-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	1
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	19
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	4
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	7
-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	-	1
-	-	-	1	1	1	1	-	-	-	-	-	-	-	-	-	1	5
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	7
-	-	-	1	-	-	-	1	-	-	-	-	-	-	-	-	3	5
-	1	1	2	2	1	5	5	1	9	1	2	7	1	9	4	28	154
-	-	1	-	1	4	1	-	1	4	-	5	2	-	6	1	276	327
-	1	2	2	3	5	6	5	2	13	1	7	9	1	15	5	304	481

Table 6
Foodborne Disease Outbreaks, by Place Where Food
Was Eaten, and Specific Etiology, 1978

	<u>Home</u>	<u>Restaurant</u>	<u>School</u>	<u>Picnic</u>	<u>Church</u>	<u>Camp</u>	<u>Other or Unknown</u>	<u>Total</u>
<u>BACTERIAL</u>								
<u>B. cereus</u>	1	3	1	-	-	-	1	6
<u>Brucella</u>	-	-	-	-	-	-	-	-
<u>C. botulinum</u>	9	3	-	-	-	-	-	12
<u>C. perfringens</u>	3	5	-	-	-	-	1	9
<u>E. coli</u>	-	-	-	-	-	1	-	1
<u>Salmonella</u>	19	15	2	1	-	-	8	45
<u>Shigella</u>	1	1	-	-	-	-	2	4
<u>S. aureus</u>	4	4	5	2	2	-	6	23
<u>Enterococci</u>	-	1	-	-	-	-	-	1
<u>Streptococcus Group A</u>	-	-	-	-	-	-	-	-
<u>V. cholerae 01</u>	1	-	-	-	-	-	-	1
<u>V. parahaemolyticus</u>	-	-	-	-	1	-	1	2
<u>Other bacteria</u>	1	-	-	-	-	-	-	1
<u>Total</u>	<u>39</u>	<u>32</u>	<u>8</u>	<u>3</u>	<u>3</u>	<u>1</u>	<u>19</u>	<u>105</u>
<u>CHEMICAL</u>								
Heavy metals	-	-	-	1	-	-	-	1
Ciguatoxin	15	3	-	-	-	-	1	19
Neurotoxic shellfish	-	-	-	-	-	-	-	-
Paralytic shellfish	3	-	-	-	-	-	1	4
Serombrotoxin	1	3	-	-	-	-	3	7
Monosodium glutamate	-	-	-	-	-	-	-	-
Mushroom poisoning	1	-	-	-	-	-	-	1
<u>Other chemicals</u>	<u>1</u>	<u>1</u>	<u>-</u>	<u>1</u>	<u>-</u>	<u>-</u>	<u>2</u>	<u>5</u>
<u>Total</u>	<u>21</u>	<u>7</u>	<u>-</u>	<u>2</u>	<u>-</u>	<u>-</u>	<u>7</u>	<u>37</u>
<u>PARASITIC</u>								
<u>T. spiralis</u>	<u>6</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>1</u>	<u>7</u>
<u>Total</u>	<u>6</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>1</u>	<u>7</u>
<u>VIRAL</u>								
Hepatitis A	1	3	-	-	-	-	1	5
<u>Total</u>	<u>1</u>	<u>3</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>-</u>	<u>1</u>	<u>5</u>
CONFIRMED TOTAL	67	42	8	5	2	1	29	154
UNKNOWN	55	192	10	4	1	6	59	327
TOTAL 1978	122	234	18	9	3	7	88	481

Table 7
Foodborne Disease Outbreaks, by Place Where Food Was Mishandled,
and Specific Etiology, 1978

	<u>Food Processing Establishments</u>	<u>Food Service Establishments</u>	<u>Homes</u>	<u>Unknown</u>	<u>Not Applicable</u>	<u>Total</u>
<u>BACTERIAL</u>						
<u>B. cereus</u>	-	4	1	1	-	6
<u>Brucella</u>	-	-	-	-	-	-
<u>C. botulinum</u>	-	1	6	5	-	12
<u>C. perfringens</u>	-	7	2	-	-	9
<u>E. coli</u>	-	1	-	-	-	1
<u>Salmonella</u>	-	19	5	21	-	45
<u>Shigella</u>	-	-	1	3	-	4
<u>S. aureus</u>	-	13	3	7	-	23
<u>Enterococci</u>	-	1	-	-	-	1
<u>Streptococcus Group A</u>	-	-	-	-	-	-
<u>V. cholerae 01</u>	-	-	1	-	-	1
<u>V. parahaemolyticus</u>	-	-	-	2	-	2
<u>Other bacterial</u>	-	-	-	1	-	1
Total	-	<u>46</u>	<u>19</u>	<u>40</u>	-	<u>105</u>
<u>CHEMICAL</u>						
Heavy metals	-	-	1	-	-	1
Ciguatoxin	-	-	-	-	19	19
Neurotoxic shellfish	-	-	-	-	-	-
Paralytic shellfish	-	-	-	-	4	4
Scrombrotoxin	-	2	-	5	-	7
Monosodium glutamate	-	-	-	-	-	-
Mushroom poisoning	-	-	-	-	1	1
<u>Other chemicals</u>	-	<u>4</u>	<u>1</u>	-	-	<u>5</u>
Total	-	<u>6</u>	<u>2</u>	<u>5</u>	<u>24</u>	<u>37</u>
<u>PARASITIC</u>						
<u>T. spiralis</u>	-	-	-	-	7	7
Total	-	-	-	-	<u>7</u>	<u>7</u>
<u>VIRAL</u>						
Hepatitis A	-	<u>2</u>	-	<u>3</u>	-	<u>5</u>
Total	-	<u>2</u>	-	<u>3</u>	-	<u>5</u>
CONFIRMED TOTAL	-	54	21	48	31	154
UNKNOWN	-	82	17	228	-	327
TOTAL 1978	-	136	38	276	31	481

Table 8
Foodborne Disease Outbreaks, by Contributing Factors, and Etiology, 1978

	Number of Reported Outbreaks	Number of Outbreaks in Which Factors Reported	Improper Holding Tempera- tures	Inade- quate Cooking	Contami- nated Equipment	Food From Unsafe Source	Poor Per- sonal Hygiene	Other
<u>BACTERIAL</u>								
<u>B. cereus</u>	6	4	4	1	1	-	-	-
<u>Brucella</u>	-	-	-	-	-	-	-	-
<u>C. botulinum</u>	12	7	-	7	-	-	-	1
<u>C. perfringens</u>	9	8	8	4	2	-	1	3
<u>E. coli</u>	1	1	1	1	1	-	-	-
<u>Salmonella</u>	45	23	16	12	11	3	9	4
<u>Shigella</u>	4	1	1	-	-	-	1	-
<u>S. aureus</u>	23	15	15	2	3	1	6	2
<u>Enterococci</u>	1	-	-	-	-	-	-	-
<u>Streptococcus Group A</u>	-	-	-	-	-	-	-	-
<u>V. cholerae O1</u>	1	1	1	1	-	-	-	-
<u>V. parahaemolyticus</u>	2	1	1	-	-	-	-	1
<u>Other bacterial</u>	1	-	-	-	-	-	-	-
Total	105	61	47	28	18	4	17	11
<u>CHEMICAL</u>								
Heavy Metal	1	-	-	-	-	-	-	-
Ciguatoxin	19	-	-	-	-	-	-	-
Neurotoxic shellfish	-	-	-	-	-	-	-	-
Paralytic shellfish	4	4	-	-	-	4	-	1
Scrombrotoxin	7	2	2	-	-	-	-	-
Monosodium glutamate	-	-	-	-	-	-	-	-
Mushroom poisoning	1	-	-	-	-	-	-	-
Other chemicals	5	2	1	-	1	-	1	1
Total	37	8	3	-	1	4	1	2
<u>PARASITIC</u>								
<u>T. spiralis</u>	7	3	-	3	-	-	-	-
Total	7	3	-	3	-	-	-	-
<u>VIRAL</u>								
Hepatitis A	5	2	-	-	-	-	2	-
Total	5	2	-	-	-	-	2	-
CONFIRMED TOTAL	154	74	50	31	19	8	20	13
UNKNOWN	327	132	100	22	26	8	43	19
TOTAL 1978	481	206	150	53	45	16	63	32

Table 9
Foodborne Disease Outbreaks, by Month of Occurrence,
and Specific Etiology, 1978

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Unknown	Total
<u>BACTERIAL</u>														
<u>B. cereus</u>	-	-	-	-	-	1	1	1	-	-	3	-	-	6
<u>Brucella</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>C. botulinum</u>	2	1	-	2	1	1	-	1	-	2	2	-	-	12
<u>C. perfringens</u>	-	1	1	-	1	1	-	1	1	1	-	2	-	9
<u>E. coli</u>	-	-	-	-	-	-	-	-	1	-	-	-	-	1
<u>Salmonella</u>	1	1	2	3	2	5	4	7	4	6	4	4	2	45
<u>Shigella</u>	-	-	-	-	-	-	-	-	-	-	-	2	2	4
<u>S. aureus</u>	-	-	3	-	-	2	1	6	1	2	6	2	-	23
<u>Enterococci</u>	-	-	-	-	-	-	-	-	-	-	-	1	-	1
<u>Streptococcus Group A</u>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<u>V. cholerae O1</u>	-	-	-	-	-	-	-	-	1	-	-	-	-	1
<u>V. parahaemolyticus</u>	-	-	-	1	-	1	-	-	-	-	-	-	-	2
<u>Other bacterial</u>	1	-	-	-	-	-	-	-	-	-	-	-	-	1
<u>Total</u>	4	3	6	6	4	11	6	16	8	11	15	11	4	105
<u>CHEMICAL</u>														
Heavy metals	-	-	-	-	-	1	-	-	-	-	-	-	-	1
Ciguatoxin	5	1	-	1	-	-	4	-	3	-	1	4	-	19
Neurotoxic shellfish	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Paralytic shellfish	-	-	-	-	-	-	-	-	3	1	-	-	-	4
Scrombrotoxin	1	-	3	-	2	-	-	-	1	-	-	-	-	7
Monosodium glutamate	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Mushroom poisoning	1	-	-	-	-	-	-	-	-	-	-	-	-	1
<u>Other chemicals</u>	1	-	-	1	-	1	-	-	1	1	-	-	-	5
<u>Total</u>	8	1	3	2	2	2	4	-	8	2	1	4	-	37
<u>PARASITIC</u>														
<u>T. spiralis</u>	-	-	-	-	2	-	1	-	1	1	-	1	1	7
<u>Total</u>	-	-	-	-	2	-	1	-	1	1	-	1	1	7
<u>VIRAL</u>														
Hepatitis A	-	-	-	-	2	-	-	1	-	-	2	-	-	5
<u>Total</u>	-	-	-	-	2	-	-	1	-	-	2	-	-	5
CONFIRMED TOTAL	12	4	9	8	10	13	11	17	17	14	18	16	5	154
UNKNOWN	24	21	26	42	32	33	21	21	21	12	29	27	18	327
TOTAL 1978	36	25	35	50	42	46	32	38	38	26	47	43	23	481

F. LINE LISTING OF FOODBORNE DISEASE OUTBREAKS

1978

F. LINE LISTING OF FOODBORNE DISEASE OUTBREAKS, 1978

Etiology	State	Number of Cases	Date of Onset	Lab Data			Location Where Food Mishandled* and Eaten
				Patient	Vehicle	Food-handler	
<u>BACTERIAL</u>							
<u>BACILLUS CEREUS</u>							
<u>B. cereus</u>	Alabama	8	11/08		+		(B) Restaurant
<u>B. cereus</u>	California	11	11/25				(B) Restaurant
<u>B. cereus</u>	Connecticut	4	7/12		+	+	(B) Office
<u>B. cereus</u>	Connecticut	12	8/15	+	+		(B) Restaurant
<u>B. cereus</u>	New Mexico	4	6/02		+		(C) Home
<u>B. cereus</u>	North Carolina	209	11/21	+	+		(D) School
<u>CLOSTRIDIUM BOTULINUM</u>							
<u>C. botulinum</u>	California	1	6/?		+		(C) Home
<u>C. botulinum</u>	California	2	10/31		+		(C) Home
<u>C. botulinum</u>	Colorado	1	1/?	+			(D) Home
<u>C. botulinum</u>	Colorado	8	11/13		+		(D) Restaurant
<u>C. botulinum</u>	Georgia	1	5/21				(D) Home
<u>C. botulinum</u>	New Mexico	34	4/09		+		(D) Restaurant
<u>C. botulinum</u>	Pennsylvania	2	4/24				(C) Home
<u>C. botulinum</u>	Texas	3	1/14				(D) Home
<u>C. botulinum</u>	Texas	1	2/06				(C) Home
<u>C. botulinum</u>	Virginia	1	10/23				(C) Home
<u>C. botulinum</u>	Wisconsin	1	11/09				(C) Home
<u>C. botulinum</u>	Puerto Rico	3	8/04				(B) Restaurant
<u>CLOSTRIDIUM PERFRINGENS</u>							
<u>C. perfringens</u>	California	18	8/19		+		(C) Home
<u>C. perfringens</u>	Georgia	2	9/14		+		(B) Restaurant
<u>C. perfringens</u>	Georgia	2	12/23	+	+	+	(B) Home
<u>C. perfringens</u>	North Carolina	16	2/22	+	+		(B) Sorority Hall
<u>C. perfringens</u>	Utah	50	3/20	+	+		(B) Restaurant
<u>C. perfringens</u>	Washington	500	5/14	+			(B) Restaurant
<u>C. perfringens</u>	Washington	5	10/02	+			(B) Restaurant
<u>C. perfringens</u>	Washington	4	12/31	+			(B) Restaurant
<u>C. perfringens</u>	Wisconsin	20	6/18		+		(C) Home
<u>ESCHERICHIA COLI</u>							
<u>E. coli</u>	Oregon	35	9/17	+	+		(B) Camp
<u>SALMONELLA</u>							
<u>S. typhimurium</u>	Arizona	23	10/?				(D) Home
<u>S. enteritidis</u>	Arkansas	7	5/14	+	+	+	(C) Home
<u>S. cubana</u>	California	5	3/06	+	+		(D) Drive In
<u>S. cubana</u>	California	13	3/12	+	+	+	(B) Home
<u>S. heidelberg</u>	California	55	7/02	+		+	(B) Other
<u>S. enteritidis</u>	California	58	7/19	+	+	+	(D) Restaurant
<u>S. schwarzengrund</u>	California	9	8/20	+	+	+	(B) Home
<u>S. typhimurium</u>	California	34	11/20	+			(B) Club Lodge
<u>S. typhimurium</u>	Colorado	7	1/18				(D) Other
<u>S. typhimurium</u>	Connecticut	11	5/30	+	+		(C) Home

*(A)--Food processing establishment; (B)--Food service establishment; (C)--Home; (D)--Unknown; (E)--Not applicable

Etiology	State	Number of Cases	Date of Onset	Lab Data			Vehicle	Location Where Food Mishandled* and Eaten
				Patient	Vehicle	Food-Handler		
<u>S. montevideo</u>	Georgia	3	4/30	+		+	Mexican food	(D) Restaurant
<u>S. derby</u>	Georgia	7	8/18	+	+		Pork	(B) Home
<u>S. javiana</u>	Georgia	756	11/05	+	+	+	Bar-B-Q pork	(B) School
<u>S. derby</u>	Georgia	7	12/17	+			Unknown	(B) Restaurant
<u>S. javiana</u>	Georgia	2	12/23	+			Pork	(B) Restaurant
<u>S. agona</u>	Kansas	6	8/05	+			Mexican food	(D) Restaurant
<u>S. heidelberg</u>	Minnesota	17	9/24		+		Hamburger casserole	(B) Home
<u>S. agona</u>	Minnesota	34	12/?		+	+	Unknown	(D) Restaurant
<u>S. typhimurium</u>	Nebraska	6	8/09		+		Chicken salad	(D) Restaurant
<u>S. enteritidis</u>	Nevada	3	6/?	+			Unknown	(D) Unknown
<u>S. enteritidis</u>	New Hampshire	4	10/05				Unknown	(D) Restaurant
<u>S. typhimurium</u>	New Jersey	26	10/29	+	+	+	Goat	(C) Home
<u>S. (not defined)</u>	New Mexico	3	9/04	+			Instant milk	(B) Home
<u>S. (not defined)</u>	New Mexico	5	?/?	+			Unknown	(D) Restaurant
<u>S. (not defined)</u>	New Mexico	5	?/?				Pork	(D) Home
<u>S. (not defined)</u>	Oklahoma	8	6/25	+			Unknown	(C) Home
<u>S. enteritidis</u>	Oregon	4	8/25				Unknown	(D) Restaurant
<u>S. typhimurium</u>	Pennsylvania	5	4/06	+	+		Egg nog	(D) Home
<u>S. typhimurium</u>	Pennsylvania	46	6/25	+			Beef	(D) Home
<u>S. typhimurium</u>	Pennsylvania	50	6/26	+			Unknown	(D) Home
<u>S. heidelberg</u>	Pennsylvania	8	7/13	+			Turkey	(D) Restaurant
<u>S. typhimurium</u>	Pennsylvania	13	9/11	+			Unknown	(D) Home
<u>S. enteritidis</u>	Tennessee	200	2/14	+		+	Turkey	(D) Club Lodge
<u>S. typhimurium</u>	Virginia	7	7/30	+	+		Ice cream	(C) Home

<u>S. typhimurium</u>	Virginia	28	8/13	+	+	+	Unknown	(B) Home
<u>S. typhimurium</u>	Virginia	110	11/12	+	+		Turkey	(B) Natl Gd Armory
<u>S. muenster</u>	Washington	27	10/03	+		+	Turkey sandwiches	(B) Restaurant
<u>S. typhimurium</u>	Wisconsin	108	8/06	+			Potato salad	(B) Picnic
<u>S. reading</u>	Wisconsin	137	11/22	+			Potatoes & gravy	(B) School
<u>S. enteritidis</u>	Wisconsin	11	12/15				Unknown	(B) Home
<u>S. typhimurium</u>	Guam	29	9/?	+			Unknown	(D) Wedding
<u>S. (not defined)</u>	New York City	12	4/23				Unknown	(D) Home
<u>S. muenster</u>	New York City	6	6/19				Unknown	(B) Restaurant
<u>S. anatum</u>	New York City	4	10/05		+		Beef	(B) Restaurant
<u>S. montevideo</u>	New York City	2	10/23	+	+	+	Beef	(B) Restaurant

SHIGELLA

<u>S. sonnei</u>	Illinois	117	12/15	+			Potato salad	(C) Banquet hall
<u>S. sonnei</u>	Illinois	36	12/28	+			Unknown	(D) Banquet hall
<u>S. sonnei</u>	New Mexico	2	?/?				Unknown	(D) Restaurant
<u>S. sonnei</u>	New Mexico	4	?/?				Chicken salad	(D) Home

STAPHYLOCOCCUS

<u>S. aureus</u>	California	28	12/22	+	+		Rice-shrimp-bacon dish	(D) Pot luck dinner
<u>S. aureus</u>	Connecticut	13	10/03		+	+	Egg salad	(D) Hospital
<u>S. aureus</u>	Florida	54	11/23		+		Ham	(B) School
<u>S. aureus</u>	Georgia	50	6/24		+		Ham	(D) Picnic
<u>S. aureus</u>	Georgia	215	6/27			+	Chicken salad	(B) School

* (A)--Food processing establishment; (B)--Food service establishment; (C)--Home; (D)--Unknown; (E)--Not applicable

Etiology	State	Number of Cases	Date of Onset	Lab Data			Vehicle	Location Where Food Mishandled* and Eaten
				Patient	Vehicle	Food-Handler		
<u>S. aureus</u>	Georgia	4	8/10		+		Ham	(B) Home
<u>S. aureus</u>	Georgia	13	9/01		+		Ham	(B) School
<u>S. aureus</u>	Kansas	2	10/16		+	+	Unknown	(D) Restaurant
<u>S. aureus</u>	Kentucky	161	11/27	+		+	Chicken salad	(B) Sr Citizen Ctr
<u>S. aureus</u>	Maryland	21	8/06		+		Ham	(B) Restaurant
<u>S. aureus</u>	Missouri	6	11/12		+		Ham	(C) Home
<u>S. aureus</u>	New York	104	8/29		+		Tuna salad, macaroni salad	(B) Jail
<u>S. aureus</u>	New York	116	11/10	+	+		Turkey salad	(B) School
<u>S. aureus</u>	North Carolina	257	11/16		+		Chicken salad	(B) Prison
<u>S. aureus</u>	Oklahoma	118	3/03	+	+	+	Potato salad, carrot salad	(B) School
<u>S. aureus</u>	Pennsylvania	29	8/25		+		Ham	(D) Church
<u>S. aureus</u>	Pennsylvania	11	8/26		+		Potato salad	(B) Church
<u>S. aureus</u>	Pennsylvania	24	11/11	+			Unknown	(D) Club lodge
<u>S. aureus</u>	South Carolina	58	7/30	+	+	+	Ham	(B) Picnic
<u>S. aureus</u>	Washington	2	3/06		+		Crab salad	(D) Restaurant
<u>S. aureus</u>	Washington	5	8/13	+	+		Ham	(B) Restaurant
<u>S. aureus</u>	Wisconsin	2	3/26		+		Ham	(C) Home
<u>S. aureus</u>	New York City	25	12/?		+		Chicken & rice	(C) Home

STREPTOCOCCUS

Enterococci	California	5	12/06				Mexican food	(B) Restaurant
-------------	------------	---	-------	--	--	--	--------------	----------------

VIBRIOS

<u>V. cholerae</u> O1	Louisiana	11	9/?			+	Crabs	(C) Home
<u>V. parahaemolyticus</u>	Louisiana	82	6/21	+	+		Shrimp	(D) Other
<u>V. parahaemolyticus</u>	Guam	4	4/07				Shrimp	(D) Other

OTHER BACTERIAL

	Utah	8	1/?	+	+		Unknown	(D) Home
--	------	---	-----	---	---	--	---------	----------

CHEMICAL

Metal	Missouri	41	6/15			+	Peach Koolaid	(C) Picnic
Ciguatoxin	Florida	3	1/05				Grouper	(E) Home
Ciguatoxin	Florida	3	7/20	+			Barracuda	(E) Restaurant
Ciguatoxin	Hawaii	2	1/09				Amberjack	(E) Home
Ciguatoxin	Hawaii	5	1/31				Amberjack	(E) Home
Ciguatoxin	Hawaii	2	4/30				Amberjack	(E) Home
Ciguatoxin	Hawaii	2	7/04				Goat fish	(E) Home
Ciguatoxin	Hawaii	3	7/16			+	Goat fish	(E) Home
Ciguatoxin	Hawaii	4	7/29				Amberjack	(E) Other
Ciguatoxin	Hawaii	2	9/20			+	Mullet	(E) Home
Ciguatoxin	Hawaii	1	9/22				Goat fish	(E) Home
Ciguatoxin	Hawaii	2	9/25			+	Gray snapper	(E) Home
Ciguatoxin	Hawaii	3	11/20			+	Goat fish	(E) Home
Ciguatoxin	Hawaii	5	12/12				Amberjack	(E) Home
Ciguatoxin	Hawaii	2	12/26			+	Porgy	(E) Restaurant
Ciguatoxin	Hawaii	2	12/29				Snapper	(E) Home

*(A)--Food processing establishment; (B)--Food service establishment; (C)--Home; (D)--Unknown; (E)--Not applicable

Etiology	State	Number of Cases	Date of Onset	Lab Data			Vehicle	Location Where Food Mishandled* and Eaten
				Patient	Vehicle	Food-Handler		
Ciguatoxin	Virgin Islands	7	1/04				Red snapper	(E) Restaurant
Ciguatoxin	Virgin Islands	1	1/15		+		Grouper	(E) Home
Ciguatoxin	Virgin Islands	6	2/11		+		Grouper	(E) Home
Ciguatoxin	Puerto Rico	1	12/06		+		Snapper	(E) Home
Paralytic shellfish	Washington	2	9/20				Mussels	(E) Home
Paralytic shellfish	Washington	3	9/23		+		Mussels	(E) Home
Paralytic shellfish	Washington	1	9/26				Mussels	(E) Home
Paralytic shellfish	Washington	4	10/03		+		Scallops	(E) Lab
Scombrotoxin	Hawaii	2	1/27				Mahi-Mahi	(B) Restaurant
Scombrotoxin	Hawaii	2	3/01		+		Mahi-Mahi	(D) Home
Scombrotoxin	Hawaii	13	3/10		+		Mahi-Mahi	(D) Hospital
Scombrotoxin	Hawaii	4	3/17				Mahi-Mahi	(D) Hospital
Scombrotoxin	Hawaii	1	5/10				Mahi-Mahi	(D) Hotel
Scombrotoxin	Hawaii	1	5/16				Mahi-Mahi	(D) Restaurant
Scombrotoxin	Hawaii	7	9/03				Mahi-Mahi	(B) Restaurant
Mushroom poisoning	California	7	1/?	+	+	+	Mushrooms	(E) Home
Other Chemical	California	2	10/04		+		Unknown	(B) Restaurant
Other Chemical	Colorado	9	4/27		+		Choc. bundt cake	(C) Office party
Other Chemical	New Jersey	2	9/10		+		Popsicle	(B) Home
Other Chemical	New Mexico	4	6/04		+		Lemonade	(B) Picnic
Other Chemical	New York City	2	1/18				Fruit cup	(B) Office

26

PARASITICTRICHINELLA SPIRALIS

<u>T. spiralis</u>	Alaska	23	???				Bear meat	(E) Home
<u>T. spiralis</u>	California	2	9/24				Pork	(E) Camping trip
<u>T. spiralis</u>	California	2	10/02		+		Bar-B-Q spare ribs	(E) Home
<u>T. spiralis</u>	Connecticut	2	7/18		+		Beef	(E) Home
<u>T. spiralis</u>	New York	2	5/14		+		Beef	(E) Home
<u>T. spiralis</u>	Rhode Island	2	12/23		+		Sausage	(E) Home
<u>T. spiralis</u>	New York City	2	5/12		+		Beef	(E) Home

VIRAL

Hepatitis A	Arkansas	82	11/?			+	Unknown	(D) Restaurant
Hepatitis A	California	36	5/?			+	Orange juice	(D) Home
Hepatitis A	Louisiana	33	5/19				Unknown	(B) Restaurant
Hepatitis A	Minnesota	91	8/?	+	+		Potato salad	(B) Country club
Hepatitis A	Texas	58	11/?			+	Unknown	(D) Restaurant

27

UNKNOWN

	Arkansas	23	4/27				Bar-B-Q Chicken/Pork	(B) Restaurant
	Arkansas	88	8/21	+			Unknown	(D) School
	California	3	1/14				Unknown	(D) Home
	California	6	1/24				Unknown	(D) Restaurant
	California	2	2/19				Clams	(C) Home
	California	2	3/04				Unknown	(D) Home

*(A)--Food processing establishment; (B)--Food service establishment; (C)--Home; (D)--Unknown; (E)--Not applicable

Etiology	State	Number of Cases	Date of Onset	Lab Data			Vehicle	Location Where Food Mishandled* and Eaten
				Patient	Vehicle	Food-Handler		
UNKNOWN	California	3	4/12				Unknown	(D) Restaurant
	California	87	4/19				Unknown	(B) Camp
	California	25	4/24		+		Fruit salad	(D) Office
	California	2	4/25				Unknown	(C) Home
	California	15	4/30				Unknown	(D) Restaurant
	California	12	5/30	+			Ice cream	(D) Home
	California	4	6/05				Unknown	(D) Restaurant
	California	5	6/07				Unknown	(D) Home
	California	2	6/11				Unknown	(B) Restaurant
	California	17	7/21		+	+	Ham & turkey salad	(D) Office
	California	5	7/26				Unknown	(D) Home
	California	6	8/16				Unknown	(D) Restaurant
	California	7	8/19				Salad dressing	(B) Restaurant
	California	10	9/11				Pizza	(B) Restaurant
	California	8	9/19				Unknown	(D) Office
	California	40	10/20				Unknown	(D) Sr. Citizen Ctr
	California	12	11/05				Unknown	(D) Restaurant
	Colorado	15	6/05				Unknown	(D) Home
	Colorado	9	11/18				Unknown	(B) Restaurant
	Connecticut	2	2/05				Unknown	(B) Restaurant
	Connecticut	10	2/10		+		Unknown	(B) Restaurant
	Connecticut	12	3/11				Unknown	(D) Other
	Connecticut	45	4/02				Unknown	(D) Restaurant
	Connecticut	21	4/02				Unknown	(D) Restaurant

28

Connecticut	2	4/10				Chinese food	(B) Home
Connecticut	21	4/16				Unknown	(D) Home
Connecticut	4	4/24				Unknown	(D) Restaurant
Connecticut	142	5/05				Unknown	(D) School
Connecticut	23	5/16		+		Clams	(B) Restaurant
Connecticut	17	5/20				Unknown	(D) Restaurant
Connecticut	4	5/28				Clams	(D) Restaurant
Connecticut	10	5/29				Clams	(D) Picnic
Connecticut	6	5/29				Clams	(D) Picnic
Connecticut	2	7/19				Clams	(B) Other
Connecticut	4	9/05				Unknown	(D) Restaurant
Connecticut	35	9/10		+		Beef	(D) Restaurant
Connecticut	19	12/10				Unknown	(B) Restaurant
Connecticut	14	12/22				Fried rice	(D) Other
Florida	2	1/12				Unknown	(C) Home
Florida	10	3/12				Unknown	(D) Restaurant
Florida	2	3/?				Unknown	(D) Delicatessen
Florida	3	4/05				Unknown	(D) Other
Florida	8	4/08				Unknown	(D) Home
Florida	2	4/09				Unknown	(D) Unknown
Florida	11	6/19				Unknown	(D) Restaurant
Florida	11	6/20				Unknown	(D) Restaurant
Florida	2	10/21				Unknown	(D) Picnic
Florida	8	10/30				Unknown	(D) Restaurant

29

*(A)--Food processing establishment; (B)--Food service establishment; (C)--Home; (D)--Unknown; (E)--Not applicable

Etiology	State	Number of Cases	Date of Onset	Lab Data			Vehicle	Location Where Food Mishandled* and Eaten
				Patient	Vehicle	Food-Handler		
UNKNOWN	Georgia	60	1/15				Unknown	(B) Restaurant
	Georgia	200	9/25				Unknown	(B) Other
	Hawaii	17	4/28				Seafood newberg	(B) Restaurant
	Hawaii	15	11/28				Unknown	(D) Other
	Idaho	35	8/07				Unknown	(D) Unknown
	Idaho	5	8/09				Unspecified meat	(B) Restaurant
	Idaho	12	8/12				Unknown	(D) Camp
	Idaho	24	8/13				Unknown	(D) Restaurant
	Illinois	74	8/05		+		Unspecified fish	(C) Home
	Illinois	21	11/30	+			Unknown	(B) Restaurant
	Indiana	74	8/15				Ham	(C) Home
	Kansas	9	4/24				Unknown	(D) Other
	Kansas	5	6/18				Unknown	(D) Home
	Kansas	14	6/25				Unknown	(B) Other
	Kansas	3	11/20				Unknown	(D) Other
	Maine	27	2/17				Unknown	(D) Other
	Maine	27	3/01				Unknown	(D) Unknown
	Maine	39	6/04				Unknown	(D) Home
	Maryland	24	3/24				Unknown	(B) Other
	Maryland	150	4/29				Unknown	(D) Restaurant
	Maryland	25	4/?				Unknown	(D) Restaurant
	Maryland	3	5/17				Unknown	(D) Home
	Maryland	135	6/13				Unknown	(B) Other
	Maryland	6	7/08				Unknown	(C) Home

	Maryland	50	7/18	+			Unknown	(B) Camp
	Maryland	2	8/15				Unknown	(B) Restaurant
	Maryland	15	9/10				Unknown	(B) Restaurant
	Maryland	42	10/28				Unknown	(B) Other
	Maryland	2	11/06				Unknown	(D) Restaurant
	Maryland	100	11/23				Turkey & dressing	(B) Restaurant
	Maryland	2	11/25				Unknown	(D) Restaurant
	Maryland	38	12/05				Unknown	(D) Restaurant
	Maryland	29	12/13				Unknown	(B) Restaurant
	Maryland	14	12/15				Unknown	(D) School
	Maryland	57	12/23	+			Unknown	(B) Restaurant
	Michigan	9	2/26	+	+		Beef	(B) Restaurant
	Michigan	110	4/21				Unknown	(B) School
	Minnesota	6	3/26				Unknown	(D) Home
	Minnesota	22	3/31				Cornish hens/dressing	(B) Restaurant
	Minnesota	2	5/18		+		Unknown	(B) Other
	Minnesota	2	7/09				Chinese food	(C) Home
	Minnesota	16	8/22		+		Unspec. baked food	(C) Home
	Minnesota	5	8/25				Chinese food	(B) Restaurant
	Minnesota	3	9/22				Unknown	(B) Restaurant
	Minnesota	15	10/26				Mexican food	(B) Restaurant
	Minnesota	9	10/27				Sausage	(B) Restaurant
	Minnesota	2	12/21				Unknown	(D) Restaurant
	Missouri	31	3/16				Unknown	(D) Restaurant

*(A)--Food processing establishment; (B)--Food service establishment; (C)--Home; (D)--Unknown; (E)--Not applicable

Etiology	State	Number of Cases	Date of Onset	Lab Data			Vehicle	Location Where Food Mishandled* and Eaten
				Patient	Vehicle	Food-Handler		
UNKNOWN	Missouri	47	6/17				Unknown	(D) Nursing home
	Missouri	8	11/14				Seafood platter	(D) Restaurant
	Montana	45	5/06				Beef stroganoff	(B) Restaurant
	Montana	40	5/14				Roast beef/sauce	(D) Restaurant
	Montana	3	12/01				Unknown	(C) Home
	Montana	27	12/15				Pork	(B) Restaurant
	Nebraska	11	7/16				Unknown	(D) Restaurant
	Nebraska	95	9/27				Unknown	(D) Restaurant
	Nebraska	8	11/18				Unknown	(D) Clinic
	Nebraska	20	12/05	+			Unknown	(D) Home
	New Hampshire	19	7/22				Unknown	(D) Wedding
	New Jersey	8	1/27				Unknown	(B) Restaurant
	New Jersey	10	3/05			+	Unknown	(D) Home
	New Jersey	74	4/23	+			Unknown	(B) Police academy
	New Jersey	12	4/24				Beef	(D) Home
	New Jersey	26	5/07				Clams	(C) Home
	New Jersey	900	5/20				Unknown	(B) Church
	New Jersey	47	6/12				Unknown	(B) Restaurant
	New Jersey	26	12/16				Unknown	(D) Restaurant
	New Mexico	4	6/29				Unknown	(C) Home
	New Mexico	2	8/26				Unknown	(C) Home
	New Mexico	2	11/08				Unknown	(B) Restaurant
	New Mexico	2	???				Unknown	(D) Unknown
	New Mexico	2	???				Unknown	(D) Restaurant

	New Mexico	2	???				Unknown	(D) Other
	New Mexico	2	???				Unknown	(D) Home
	New Mexico	19	???				Beef	(D) Restaurant
	New Mexico	50	???				Other, not spec	(D) School
	New Mexico	3	???				Chicken	(D) Restaurant
	New Mexico	5	???				Unspecified diary	(D) Restaurant
	New Mexico	4	???				Chicken	(D) Home
	New Mexico	2	???				Chicken	(D) Home
	New Mexico	3	???				Unknown	(D) Restaurant
	New Mexico	4	???				Unspec salads/sauce	(D) Restaurant
	New Mexico	2	???				Chicken salad	(D) Restaurant
	New York	18	8/14				Ham	(D) Home
	North Carolina	8	4/18				Unknown	(D) Showroom
	Oklahoma	40	???				Unknown	(D) Other
	Oregon	7	3/27				Unknown	(D) Restaurant
	Oregon	17	6/11				Unknown	(D) Other
	Oregon	27	7/22				Unknown	(D) Camp
	Oregon	14	8/03				Unknown	(D) Home
	Oregon	5	8/30			+	Mexican food	(B) Restaurant
	Pennsylvania	4	1/18				Unknown	(D) Home
	Pennsylvania	3	3/23				Unknown	(D) Pizzeria
	Pennsylvania	2	4/01				Unknown	(D) Unknown
	Pennsylvania	15	4/03				Unknown	(D) Private club
	Pennsylvania	19	4/30				Shrimp	(D) Restaurant

*(A)--Food processing establishment; (B)--Food service establishment; (C)--Home; (D)--Unknown; (E)--Not applicable

Etiology	State	Number of Cases	Date of Onset	Lab. Data			Vehicle	Location Where Food Mishandled* and Eaten
				Patient	Vehicle	Food-Handler		
UNKNOWN	Pennsylvania	4	5/01				Pizza	(D) Home
	Pennsylvania	2	5/19				Unknown	(D) Restaurant
	Pennsylvania	3	5/20				Unknown	(D) Restaurant
	Pennsylvania	83	5/26				Unknown	(B) Camp
	Pennsylvania	34	6/05				Cake	(D) Picnic
	Pennsylvania	9	6/18				Unknown	(B) Camp
	Pennsylvania	61	7/20				Egg salad	(D) Other
	Pennsylvania	4	7/22				Unknown	(D) Home
	Pennsylvania	6	7/30				Unknown	(D) Home
	Pennsylvania	25	10/24				Unknown	(B) Cafeteria
	Pennsylvania	5	11/16				Unknown	(B) Home
	Pennsylvania	15	12/04				Unknown	(D) Banquet hall
	Rhode Island	80	1/05				Turkey & gravy	(D) School
	South Carolina	3	6/19				Unknown	(B) Break room
	South Dakota	16	5/22				Unknown	(D) Mil. kitchen
	Texas	45	5/05				Unknown	(D) School
	Vermont	44	2/05				Unknown	(D) Restaurant
	Virginia	3	3/?				Unknown	(D) Office
	Virginia	18	4/16		+		Beef	(D) Fast food
	Virginia	50	5/16				Unknown	(B) Restaurant
	Virginia	10	10/13				Unknown	(B) Office
	Washington	3	1/29				Unknown	(B) Restaurant
	Washington	3	2/02				Unknown	(D) Restaurant
	Washington	2	2/05				Unknown	(B) Restaurant

	Washington	2	2/07				Unknown	(B) Restaurant
	Washington	2	3/05				Unknown	(D) Restaurant
	Washington	53	3/17		+		Unknown	(B) Sr citizen ctr
	Washington	5	3/23				Unknown	(B) Restaurant
	Washington	2	3/23				Unknown	(B) Restaurant
	Washington	10	4/04				Unknown	(B) Restaurant
	Washington	2	4/06				Unknown	(B) Restaurant
	Washington	2	4/07				Unknown	(B) Restaurant
	Washington	2	4/16		+		Unknown	(B) Restaurant
	Washington	4	4/23				Unknown	(B) Restaurant
	Washington	3	4/23				Unknown	(D) Home
	Washington	2	5/02				Unknown	(D) Home
	Washington	2	5/06				Unknown	(B) Restaurant
	Washington	3	6/01				Unknown	(B) Restaurant
	Washington	2	6/01				Unknown	(D) Restaurant
	Washington	3	6/12				Unknown	(B) Restaurant
	Washington	14	6/14				Unknown	(B) Restaurant
	Washington	2	6/18				Unknown	(B) Restaurant
	Washington	6	6/20				Unknown	(D) Restaurant
	Washington	4	7/07				Fried rice	(B) Restaurant
	Washington	3	7/11				Unknown	(B) Restaurant
	Washington	2	7/13		+		Unknown	(B) Restaurant
	Washington	4	7/15				Unknown	(B) Restaurant
	Washington	2	7/29				Chinese food	(B) Restaurant

*(A)--Food processing establishment; (B)--Food service establishment; (C)--Home; (D)--Unknown; (E)--Not applicable

Etiology	State	Number of Cases	Date of Onset	Lab Data			Vehicle	Location Where Food Mishandled* and Eaten
				Patient	Vehicle	Food-Handler		
UNKNOWN	Washington	5	8/05				Unknown	(C) Home
	Washington	2	8/22				Unknown	(D) Restaurant
	Washington	21	9/07				Unknown	(C) School
	Washington	2	9/18				Unknown	(D) Restaurant
	Washington	2	9/24				Unknown	(D) Restaurant
	Washington	2	9/25				Unknown	(B) Restaurant
	Washington	6	11/09				Unknown	(D) Home
	Washington	4	11/09				Unknown	(D) Home
	Washington	3	11/14				Unknown	(B) Restaurant
	Washington	4	11/24				Unknown	(B) Restaurant
	Washington	2	11/27				Unknown	(D) Restaurant
	Washington	2	12/03				Unknown	(C) Home
	Washington	22	12/10				Unknown	(D) Restaurant
	Washington	54	12/13				Unknown	(B) Cafeteria
	Wisconsin	4	2/14				Unknown	(C) Home
	Wisconsin	51	12/09				Unknown	(D) Restaurant
	Guam	10	5/06		+		Shellfish	(C) Home
	Guam	23	12/09				Unknown	(B) Prison
	New York City	2	1/07				Unknown	(D) Restaurant
	New York City	2	1/07				Unknown	(D) Restaurant
	New York City	3	1/10				Unknown	(D) Other
	New York City	4	1/14				Unknown	(D) Restaurant
	New York City	2	1/16				Unknown	(D) Restaurant
	New York City	2	1/20				Unknown	(D) Other

36

New York City	2	1/21				Unknown	(D) Restaurant
New York City	67	1/22				Unknown	(D) Other
New York City	4	1/22				Unknown	(D) Restaurant
New York City	3	1/22				Unknown	(D) Restaurant
New York City	2	1/25				Unknown	(D) Restaurant
New York City	4	1/25				Unknown	(D) Other
New York City	2	1/26				Unknown	(D) Restaurant
New York City	8	1/27				Unknown	(D) Restaurant
New York City	3	1/27				Unknown	(D) Home
New York City	3	1/31				Unknown	(D) Restaurant
New York City	2	2/02				Unknown	(D) Restaurant
New York City	3	2/02				Unknown	(D) Other
New York City	3	2/04				Unknown	(D) Home
New York City	2	2/05				Unknown	(D) Restaurant
New York City	2	2/05				Unknown	(D) Restaurant
New York City	3	2/07				Unknown	(D) Restaurant
New York City	2	2/25				Unknown	(D) Restaurant
New York City	3	2/26				Unknown	(D) Restaurant
New York City	2	2/26				Unknown	(D) Restaurant
New York City	2	2/28				Unknown	(D) Restaurant
New York City	4	2/?			+	Unknown	(D) Restaurant
New York City	3	3/01			+	Unknown	(D) Home
New York City	3	3/10				Unknown	(D) Restaurant
New York City	2	3/11				Unknown	(D) Restaurant

37

*(A)—Food processing establishment; (B)—Food service establishment; (C)—Home; (D)—Unknown; (E)—Not applicable

Etiology	State	Number of Cases	Date of Onset	Lab Data			Vehicle	Location Where Food Mishandled* and Eaten
				Patient	Vehicle	Food-Handler		
UNKNOWN	New York City	2	3/14				Unknown	(D) Restaurant
	New York City	2	3/14				Unknown	(D) Home
	New York City	4	3/18				Unknown	(D) Restaurant
	New York City	3	3/19				Unknown	(D) Restaurant
	New York City	2	3/20				Unknown	(D) Restaurant
	New York City	2	3/26				Unknown	(D) Restaurant
	New York City	4	4/02				Unknown	(D) Restaurant
	New York City	2	4/04				Unknown	(D) Restaurant
	New York City	2	4/08				Unknown	(D) Restaurant
	New York City	2	4/09				Unknown	(D) Restaurant
	New York City	2	4/09				Unknown	(D) Restaurant
	New York City	2	4/10				Unknown	(D) Restaurant
	New York City	7	4/13				Unknown	(D) Home
	New York City	3	4/18				Unknown	(D) Restaurant
	New York City	4	4/20				Unknown	(D) Other
	New York City	2	4/23				Unknown	(D) Restaurant
	New York City	5	5/02				Unknown	(D) Restaurant
	New York City	3	5/03				Unknown	(D) Home
	New York City	2	5/08				Unknown	(D) Restaurant
	New York City	2	5/09				Unknown	(D) Restaurant
	New York City	2	5/11				Unknown	(D) Restaurant
	New York City	2	5/21				Unknown	(D) Restaurant
	New York City	35	5/22				Unknown	(D) Other
	New York City	3	5/27				Unknown	(D) Restaurant

38

	New York City	2	5/31				Unknown	(D) School
	New York City	5	6/01				Unknown	(D) Restaurant
	New York City	36	6/04				Unknown	(D) Restaurant
	New York City	2	6/07				Unknown	(D) Restaurant
	New York City	4	6/08				Unknown	(D) Restaurant
	New York City	35	6/11				Unknown	(D) Home
	New York City	3	6/13				Unknown	(D) Restaurant
	New York City	30	6/22				Unknown	(B) Restaurant
	New York City	9	6/22				Unknown	(B) Other
	New York City	2	6/27				Unknown	(B) Restaurant
	New York City	2	6/28				Unknown	(D) Restaurant
	New York City	2	7/11				Unknown	(D) Restaurant
	New York City	2	7/12				Unknown	(B) Restaurant
	New York City	2	7/12				Unknown	(D) Restaurant
	New York City	2	7/22				Unknown	(D) Restaurant
	New York City	4	8/07				Unknown	(D) Restaurant
	New York City	2	8/12				Unknown	(D) Restaurant
	New York City	3	8/28				Unknown	(D) Restaurant
	New York City	2	9/05				Unknown	(D) Restaurant
	New York City	2	9/16				Unknown	(D) Restaurant
	New York City	2	9/17				Unknown	(D) Restaurant
	New York City	2	9/19				Unknown	(D) Home
	New York City	2	9/22				Unknown	(D) Restaurant
	New York City	5	9/26				Unknown	(D) Restaurant

39

*(A)--Food processing establishment; (B)--Food service establishment; (C)--Home; (D)--Unknown; (E)--Not applicable

<u>Etiology</u>	<u>State</u>	<u>Number of Cases</u>	<u>Date of Onset</u>	<u>Lab Data</u>			<u>Vehicle</u>	<u>Location Where Food Mishandled* and Eaten</u>
				<u>Patient</u>	<u>Vehicle</u>	<u>Food-Handler</u>		
UNKNOWN	New York City	4	9/26				Unknown	(D) Restaurant
	New York City	3	9/28				Unknown	(D) Restaurant
	New York City	2	9/29				Unknown	(D) Restaurant
	New York City	2	10/08				Unknown	(D) Restaurant
	New York City	2	10/11				Unknown	(D) Restaurant
	New York City	39	10/15				Unknown	(B) Restaurant
	New York City	3	10/15				Unknown	(D) Restaurant
	New York City	50	11/07				Unknown	(D) Other
	New York City	2	11/07				Unknown	(D) Other
	New York City	3	11/07				Unknown	(D) Other
	New York City	3	11/09				Unknown	(D) Restaurant
	New York City	2	11/11				Unknown	(D) Restaurant
	New York City	2	11/14				Unknown	(D) Restaurant
	New York City	3	11/19				Unknown	(D) Other
	New York City	2	11/20				Unknown	(D) Restaurant
	New York City	2	11/26				Unknown	(D) Restaurant
	New York City	2	11/26				Unknown	(D) Restaurant
	New York City	2	11/29				Unknown	(D) Restaurant
	New York City	2	11/30				Unknown	(D) Restaurant
	New York City	2	12/02				Unknown	(D) Restaurant
	New York City	12	12/02				Unknown	(D) Home
	New York City	2	12/07				Unknown	(D) Restaurant
	New York City	25	12/09				Unknown	(D) Restaurant
	New York City	2	12/11				Unknown	(D) Restaurant

	New York City	3	12/11				Unknown	(D) Restaurant
	New York City	2	12/15				Unknown	(D) Restaurant
	New York City	2	12/20				Unknown	(D) Other
	New York City	4	12/20				Unknown	(D) Restaurant
	New York City	2	12/30				Unknown	(D) Restaurant
	New York City	12	??				Unknown	(B) Other
	New York City	20	??				Unknown	(D) School
	New York City	2	??				Unknown	(D) Restaurant
	New York City	2	??				Unknown	(D) Restaurant

*(A)--Food processing establishment; (B)--Food service establishment; (C)--Home; (D)--Unknown; (E)--Not applicable

G. Guidelines for Confirmation of Foodborne Disease Outbreak

<u>BACTERIAL</u>	<u>Clinical Syndrome</u>	<u>Laboratory and/or Epidemiologic Criteria</u>
1. <u>Bacillus cereus</u>	a) incubation period 2-16 hrs. b) gastrointestinal syndrome	a) isolation of $>10^5$ organisms per gram in epidemiologically incriminated food OR b) isolation of organism from stools of ill person.
2. <u>Brucella</u>	a) incubation period several days to several months b) clinical syndrome compatible with brucellosis	a) 4-fold increase in titer OR b) positive blood culture
3. <u>Clostridium botulinum</u>	a) incubation 2 hours-8days, usually 12-48 hours b) clinical syndrome compatible with botulism (see CDC Botulism Manual)	a) detection of botulinum toxin in human sera, feces, or food OR b) isolation of <u>C. botulinum</u> organism from epidemiologically incriminated food or stools OR c) food epidemiologically incriminated
4. <u>Clostridium perfringens</u>	a) incubation period 9-15 hrs. b) lower intestinal syndrome--majority of cases with diarrhea but little vomiting or fever	a) organisms of same serotype in epidemiologically incriminated food and stool of ill individuals. OR b) isolation of organisms with same serotype in stool of most ill individuals and not in stool of controls OR c) $>10^5$ organisms per gram in epidemiologically incriminated food provided specimen properly handled
5. <u>Escherichia coli</u>	a) incubation period 6-36 hrs. b) gastrointestinal syndrome--majority of cases with diarrhea	a) demonstration of organisms of same serotype in epidemiologically incriminated food and stool of ill individuals and not in stool of controls OR b) isolation from stool of most ill individuals, organisms of the same serotype which have been shown to be enterotoxigenic or invasive by special laboratory techniques
6. <u>Salmonella</u>	a) incubation period 6-48 hrs. b) gastrointestinal syndrome--majority of cases with diarrhea	a) isolation of <u>Salmonella</u> organism from epidemiologically implicated food OR b) isolation of <u>Salmonella</u> organism from stools of ill individuals

	<u>Clinical Syndrome</u>	<u>Laboratory and/or Epidemiologic Criteria</u>
7. <u>Shigella</u>	a) incubation period 12-50 hours b) gastrointestinal syndrome-- majority of cases with diarrhea	a) isolation of <u>Shigella</u> organ- ism from epidemiologically im- plicated food <u>OR</u> b) isolation of <u>Shigella</u> organ- ism from stools of ill individ- uals
8. <u>Staphylococcus aureus</u>	a) incubation period 30 min.- 8 hours (usually 2-4 hrs.) b) gastrointestinal syndrome-- majority of cases with vomiting	a) detection of enterotoxin in epidemiologically implicated food <u>OR</u> b) organisms with same phage type in stools or vomitus of ill individuals and, when pos- sible, implicated food and/or skin or nose of food handler <u>OR</u> c) isolation of $>10^5$ organ- isms per gram in epidemiologi- cally implicated food
9. <u>Streptococcus Group A</u>	a) incubation period 1-4 days b) febrile URI syndrome	a) isolation of organisms with same M and T type from impli- cated food <u>OR</u> b) isolation of organisms with same M and T type from throats of ill individuals
10. <u>Vibrio cholerae</u>	a) incubation period 1-3 days b) gastrointestinal syndrome-- majority of cases with diarrhea and without fever	a) isolation of <u>V. cholerae</u> from epidemiologically incrim- inated food <u>OR</u> b) isolation of organisms from stools or vomitus of ill individuals <u>OR</u> c) significant rise in vibrio- cidal, bacterial agglutinating or antitoxin antibodies in acute and early convalescent sera, or significant fall in vibriocidal antibodies in early and late convalescent sera in persons not recently immunized
11. <u>Vibrio parahaemolyticus</u>	a) incubation period 15-24 hrs. b) gastrointestinal syndrome-- majority of cases with diarrhea	a) isolation of $>10^5$ organ- isms from epidemiologically implicated food (usually seafood) <u>OR</u> b) isolation of Kanagawa- positive organisms of same serotype from stool of ill individuals
12. Others	clinical data appraised in individual circumstances	laboratory data appraised in individual circumstances

Clinical Syndrome

Laboratory and/or
Epidemiologic Criteria

CHEMICAL

<p>1. Heavy metals</p> <p>Antimony Cadmium Copper Iron Tin Zinc, etc</p>	<p>a) incubation period 5 min. to 8 hrs. (usually less than 1 hr)</p> <p>b) clinical syndrome compatible with heavy metal poisoning--usually gastrointestinal syndrome and often metallic taste</p>	<p>demonstration of high concentration of metallic ion in epidemiologically incriminated food or beverage</p>
<p>2. Ichthyosarcotoxin</p>		
<p>Ciguatoxin</p>	<p>a) incubation period 1-36 hrs. (usually 2-8 hrs.)</p> <p>b) clinical syndrome compatible with ciguatera--usually initial gastrointestinal symptoms followed by dry mouth, paresthesia of lips, tongue, throat or extremities. A sensation of looseness and pain in the teeth and and a paradoxical temperature sensation are characteristic</p>	<p>a) demonstration of ciguatoxin in epidemiologically incriminated fish</p> <p style="text-align: center;"><u>OR</u></p> <p>b) ciguatera-associated fish epidemiologically incriminated</p>
<p>Puffer fish (tetrodotoxin)</p>	<p>a) incubation period 10 min. to 3 hrs. (usually 10-45 min.)</p> <p>b) clinical syndrome compatible with puffer fish poisoning--paresthesia of lips, tongue, face or extremities often followed by numbness, loss of proprioception or a "floating" sensation</p>	<p>a) demonstration of tetrodotoxin in fish</p> <p style="text-align: center;"><u>OR</u></p> <p>b) puffer fish epidemiologically incriminated</p>
<p>Scombrototoxin</p>	<p>a) incubation period 1 min. to 3 hours (usually less than 1 hour)</p> <p>b) clinical syndrome compatible with scombroid fish poisoning often including flushing, headache, dizziness, burning of mouth and throat, upper and lower gastrointestinal symptoms, urticaria and generalized pruritus</p>	<p>a) demonstration of elevated histamine levels in epidemiologically incriminated fish</p> <p style="text-align: center;"><u>OR</u></p> <p>b) fish of order Scombroidei or fish associated with scombroid poisoning (e.g., mahi-mahi) epidemiologically incriminated</p>

	<u>Clinical Syndrome</u>	<u>Laboratory and/or Epidemiologic Criteria</u>
3. Monosodium glutamate	<p>a) incubation period 3 min. to 2 hours (usually less than 1 hour)</p> <p>b) clinical syndrome compatible with monosodium glutamate intoxication--often including burning sensations in chest, neck, abdomen or extremities, sensations of lightness and pressure over face, or a heavy feeling in the chest</p>	<p>history of large amounts (usually >1.5 grams) of MSG having been added to epidemiologically incriminated food</p>
<hr/>		
4. Mushroom poison		
Group containing ibotenic acid and muscimol	<p>a) incubation period 1-12 hrs. (usually less than 4 hrs.)</p> <p>b) clinical syndrome compatible with mushroom poisoning by this group--often including confusion, delirium, visual disturbances</p>	<p>a) demonstration of toxic chemical in epidemiologically incriminated mushrooms</p> <p style="text-align: center;">OR</p> <p>b) epidemiologically incriminated mushrooms identified as a toxic type</p>
Group containing amanitotoxins and phallotoxins, or gyromitrin	<p>a) incubation period 5-18 hrs.</p> <p>b) characteristic clinical syndrome compatible with mushroom poisoning by this group--upper and lower gastrointestinal symptoms followed by hepatic and/or renal failure</p>	<p>a) demonstration of toxic chemical in epidemiologically incriminated mushrooms</p> <p style="text-align: center;">OR</p> <p>b) epidemiologically incriminated mushrooms identified as a toxic type</p>
Groups containing muscarine, psilocybin and psilocin, gastrointestinal irritants, disulfiram-like compounds	<p>a) characteristic incubation period</p> <p>b) clinical syndrome compatible with mushroom poisoning by these groups</p>	<p>a) demonstration of toxic chemical in epidemiologically incriminated mushrooms</p> <p style="text-align: center;">OR</p> <p>b) epidemiologically incriminated mushroom identified as toxic type</p>
5. Paralytic and neurotoxic shellfish poison	<p>a) incubation period 30 min. to 3 hours</p> <p>b) clinical syndrome compatible with paralytic shellfish poisoning--often including paresthesia of lips, mouth or face and often upper and lower gastrointestinal symptoms</p>	<p>a) detection of toxin in epidemiologically incriminated mollusks</p> <p style="text-align: center;">OR</p> <p>b) detection of large numbers of shellfish poisoning-associated species of dinoflagellates in water from which epidemiologically incriminated mollusks gathered</p>
6. Other chemical	clinical data appraised in individual circumstances	laboratory data appraised in individual circumstances

PARASITIC AND VIRAL

1. <u>Trichinella spiralis</u>	a) incubation period 3-30 days	a) muscle biopsy from ill individual
--------------------------------	--------------------------------	--------------------------------------

	<u>Clinical Syndrome</u>	<u>Laboratory and/or Epidemiologic Criteria</u>
	b) clinical syndrome compatible with trichinosis--often including fever, high eosinophil count, orbital edema, myalgia	<u>OR</u> b) serological tests <u>OR</u> c) demonstration of larvae in incriminated food
2. Hepatitis A	a) incubation period 10-45 days b) clinical syndrome compatible with hepatitis--usually including jaundice, GI symptoms, dark urine	liver function tests compatible with hepatitis in affected persons who consumed the epidemiologically incriminated food
3. Others	clinical evidence appraised in individual circumstances	laboratory evidence appraised in individual circumstances

H. Selected Foodborne Outbreak Articles, Taken from Morbidity and Mortality Weekly Report

Clostridium perfringens Food Poisoning--California
(MMWR 27(19):164-165, 1978)

An outbreak of Clostridium perfringens food poisoning in Ojai, California, traced to the consumption of bean-filled burritos,* illustrates that foods other than meat, poultry, or gravy contain the essential amino acids to support growth of this organism.

The burritos that caused the outbreak were one of many Mexican-style foods offered for sale at an outdoor fund-raising event on September 18, 1977, in Ojai. On September 19, the Ventura County Environmental Health Division began receiving reports of illness and initiated an investigation. To identify cases, hospital and community doctors, school nurses, and the event-organizers were contacted, and press releases were distributed to the local media. By these means, 181 ill persons who had attended the event were identified. Information about 40 other persons who ate at the event but did not become ill was also obtained. It is not known how many persons consumed burritos at this event, although sales receipts showed that about 1,200 had been sold.

Symptoms consisted primarily of diarrhea (96.2%) and cramps (79.7%). Only 3 (1.7%) of those ill reported vomiting. The mean incubation period was 11 hours, with 87% of persons reporting illness from 8-22 hours after eating. The majority of ill persons were free of symptoms within 24 hours of onset. No one was hospitalized. Analysis of food histories incriminated the bean-filled burritos ($p < .01$).

Containers of leftover bean-burrito filling, green chili sauce, taco sauce, and shredded longhorn cheese refrigerated at 5°C (40°F) were sampled by county investigators 3 hours after closing of the food stand. Samples of 1 frozen burrito, another held at room temperature, and a frozen enchilada were also obtained. All of the food samples were examined for the presence of Salmonella, Shigella, Staphylococcus, and C. perfringens and for total aerobic colony counts. The bean-burrito filling and the unrefrigerated whole burrito were found to contain 4.0×10^6 and 7.1×10^6 C. perfringens bacteria per gram, respectively. None of the other food specimens contained more than 30,000 C. perfringens organisms per gram. No Shigella, Salmonella, or Staphylococcus organisms were found in food specimens.

The bean-burrito filling contained no meat or meat extracts. It was prepared from dried pinto beans that had been boiled with water in metal pots, mashed, and stored in cafeteria refrigerators overnight. The pots of beans were kept refrigerated while they were being transported the following morning. They were transferred to smaller open containers and reheated for an undetermined period of time before being served in the burritos. Several persons throughout the day reported that the beans were not heated and the burritos were served at ambient temperature. During that day, shade temperatures reached 29.4°C (85°F). There was no evidence of cross contamination with meat products.

Editorial Note: Pinto beans and other legumes provide an excellent substrate for C. perfringens (1). If beans are not served immediately after cooking, they should be held at $>60^\circ\text{C}$ (140°F) or rapidly cooled in shallow containers in refrigerators and reheated to 74°C (165°F) before being served.

Reference

1. Rockland LB, Gardiner BL, Pieczarka D. Stimulation of gas production and growth of Clostridium perfringens type A (No. 3624) by legumes. J Food Sci 1969;34:411-414.

*burrito: a flour tortilla filled with either meat or beans and generally garnished with hot sauce and cheese

Restaurant Outbreak of Salmonellosis Due to Undercooked Turkey--Washington
(MMWR 27(51):514, 519, 1978)

During October an outbreak of febrile gastroenteritis due to Salmonella muenster, involving 19 persons, occurred in King County, Washington. Investigation traced the illness to turkey prepared at a local restaurant.

An epidemiologic investigation was initiated on October 9, when individuals representing 3 separate groups telephoned complaints of food poisoning to the county health department. Investigation and reports by physicians identified a total of 19 patients. Eighteen of these persons had eaten at 1 restaurant at different times during October 3-5, 1978; the other patient was a cook at the restaurant.

The symptoms of the 18 diners were diarrhea (100%), abdominal cramps (100%), fever (67%), and vomiting (44%) 8 to 39 hours (mean, 17 hours) after eating. The duration of illness ranged from 1 to 17 days (mean, 3 days). S. muenster was isolated from 10 stool specimens submitted by 11 of the 18 symptomatic patients, from 1 of 2 specimens submitted by asymptomatic customers, and from the cook who had had diarrheal illness during the same 3-day period.

Food histories of the 18 customers and of 9 other restaurant patrons showed that all 18 ill diners but only 2 of 8 well customers had eaten cold turkey in either cold sandwiches or turkey salad ($p=.0012$). Investigation revealed that all turkey served in the restaurant was cooked on the premises by roasting to an internal meat temperature of 135-140°F (57-60°C), rather than the 165°F (74°C) required by Washington state regulations. Cooked turkeys were then refrigerated until served in sandwiches or salad. Hot turkey dishes were prepared by reheating sliced meat to a temperature over 164°F (74°C). No food specimens served during the implicated period were available for culture; a specimen of turkey obtained a week after the outbreak was culture-negative for salmonellae, as were several environmental cultures.

The restaurant complied promptly with all regulations; subsequently, no further cases have been reported.

Editorial Note: This outbreak of S. muenster enteritis serves as a timely reminder of the need for proper cooking of poultry and other products to prevent salmonellosis. In this outbreak, reheating apparently provided sufficient additional cooking, since only cold turkey dishes were implicated. The cook, although he was culture-positive, was probably not the source of the outbreak. He ate at the restaurant and probably became infected there.

Botulism--Puerto Rico
(MMWR 27(38):356-357, 1978)

The first outbreak of botulism reported in Puerto Rico occurred in August 1978, among a restaurant owner and 2 of his employees in Guaynabo.

On August 10, a 46-year-old Guaynabo businessman presented to a hospital emergency room with vomiting, ptosis, and blurred vision. No sensory abnormalities were noted. Initial treatment included intravenous fluids and observation, but as dysphagia and dyspnea developed, a neurologist was consulted, and the clinical diagnosis of botulism was made. He was admitted to a medical center in San Juan, where he was placed on ventilatory assistance. He received 4 vials of trivalent (ABE) botulinal antitoxin within 24 hours without noticeable improvement. He died of pneumonia on August 12.

The Puerto Rico Department of Health was notified of the case on August 10 and began an investigation immediately. It was found that the patient's wife had prepared marinated fish on approximately July 26. She had stored it in 3 large, narrow-mouthed, glass jars, closed with screw caps, and left it to "cure" under a table in her husband's cafeteria-pizzeria business. When the investigators found the jars, a thick layer of oil had formed between the fish mixture and the air remaining in each jar. The jars were confiscated.

Attempts by investigators to find who else might have consumed the fish led to the identification of 2 more cases. The second patient was a 24-year-old restaurant

employee who had developed weakness on August 4 after eating fish for 3 days. He had been hospitalized on August 8 with weakness, neck pain, and blurred vision. He never developed respiratory symptoms but had ptosis, shoulder and neck weakness, and dysphagia. By August 10, when the diagnosis of botulism was made, he was slightly improved and was not given antitoxin.

Another employee, a 16-year-old man, was absent from work on August 10. He was found at home in a remote rural community with severe weakness, ptosis, and dysarthria. He had eaten small amounts of fish on August 8 and 9. His therapy included ventilatory assistance and 4 vials of antitoxin. He developed pneumonia, and has improved slowly.

Two other employees and 3 additional people ate the fish between July 27 and August 8, but they remained well.

Serum was obtained from the patients on August 10, and type A botulinum toxin was present in patients 1 and 3. Stool, obtained only from the second patient, had no toxin. Culture results are pending. The contents of the 3 jars of marinated fish all had type A toxin; pH levels ranged from 3.9 to 4.9.

Editorial Note: Marinated fish is a very common food in Puerto Rico. Perhaps the most significant factor in toxin production in this case was the use of closed, narrow-mouthed jars rather than the customary practice of marinating fish in wide, shallow trays. The thick oil layer between the fish and the air may have protected the fish from the acid marinade while preventing passage of oxygen into the solution. Although the pH level of the food approximated the 4.5 value generally regarded as inhibitory to germination of Clostridium botulinum spores (1), the variation in pH levels in the layers present may have allowed for nonprotective regions.

Although type E toxin is commonly associated with outbreaks traced to fish, type A toxin is also frequently responsible (2,3).

Prompt notification of the Puerto Rico Health Department made possible the identification of cases 2 and 3, confiscation of the implicated food, and control of the outbreak.

References

1. Ingram M, Robinson RH. A discussion of the literature on botulism in relation to acid foods. *J Appl Bacteriol* 1951;14:73-84.
2. Merston MH, Hughes JM, Dowell VR, et al. Current trends in botulism in the United States. *JAMA* 1974;229:1305.
3. Center for Disease Control. Botulism in the United States, 1899-1973. Handbook for epidemiologists, clinicians, and laboratory workers. Issued May 1977.

Follow-up on Vibrio cholerae Infection--Louisiana (MMWR 27(39):367, 1978)

A fourth person clinically ill with cholera was identified in Louisiana. The patient, a 19-year-old woman who lives near Abbeville, had onset of diarrhea on September 18, 1978. She was hospitalized on September 21, treated with tetracycline, and recovered. Isolates from this case, the other 3 clinically ill persons, and 1 asymptomatic case were confirmed as Vibrio cholerae, biotype El Tor, serotype Inaba.

The 4 clinically ill persons had a history of ingesting steamed or boiled crab in the 2- to 5-day period before onset of illness; the asymptomatic case had also recently eaten such food. For each of the 4 ill persons, 2 matched controls were questioned about eating seafood. None of these 8 controls had recently eaten steamed or boiled crab. An isolate of V. cholerae, serotype Inaba, was made from a sample of shrimp collected in the area where crabs eaten by 1 patient were collected.

Editorial Note: These 5 persons come from 2 adjacent towns and had no single common contact or water supply. The seafoods eaten by these persons came from different locations along 60 miles of the Louisiana Gulf Coast. If the steamed crabs were a vehicle, they were insufficiently cooked to destroy V. cholerae organisms or they were

recontaminated after cooking. Residents of the area were warned to take extra care with the preparation of crabs to insure that they were adequately cooked and not subsequently contaminated.

Since the United States will now be listed by the World Health Organization as having a cholera-infected area, the following countries will now require International Certificates of Vaccination against Cholera from travelers arriving from Vermilion Parish, Louisiana: Albania, Angola, Brunei, Cape Verde, China (People's Republic), China (Republic of), Egypt, Fiji, Iran, Iraq, Lao People's Democratic Republic, Libyan Arab Jamahiriya, Madagascar, Mali, Nauru, Pakistan, Panama, Pitcairn Island, Qatar, Ryukyu Islands, Saint Helena, Seychelles, Swaziland, Yemen, and Zambia. Five countries that always require cholera vaccination from all travelers are Malawi, Maldives, Mozambique, Papua New Guinea, and Saudi Arabia. The following countries will require a Certificate only from travelers proceeding to a country with a cholera requirement: Burma, India, and Nigeria.

An area is considered infected until 10 days has passed "since the last case identified has died, recovered, or been isolated, and there is no epidemiological evidence of spread of that disease to any contiguous area" (1).

Reference

1. World Health Organization: International Health Regulations (1969). 2nd ed. Geneva, World Health Organization, 1974

I. Bibliography

GENERAL

1. Riemann H (ed): Foodborne Infections and Intoxications. New York, Academic Press, 1969
2. Food Research Institute: Annual Report 1977, University of Wisconsin, Madison, Wisconsin.
3. Bryan FL: Emerging foodborne diseases. I. Their surveillance and epidemiology. II. Factors that contribute to outbreaks and their control. J Milk Food Technol 35:618-625, 632-638, 1972
4. Horwitz MA: Specific diagnosis of foodborne disease. Gastroent 73:375-381, 1977
5. Bryan FA: Factors that contribute to outbreaks of foodborne disease. J of Food Protection 41:816-827, 1978

BACTERIAL

Bacillus cereus

1. Goepfert JM, Spira WM, Kim HU: Bacillus cereus: Food poisoning organism. A review. J Milk Food Technol 35:213-227, 1972
2. Mortimer PR, McCann G: Food poisoning episodes associated with Bacillus cereus in fried rice. Lancet 1:1043-1045, 1974
3. Terranova W, Blake PA: Bacillus cereus food poisoning. N Eng J Med 298:143-144, 1978

Brucella

1. Spink WW: The Nature of Brucellosis. Minneapolis, Lund Press, Inc., 1956
2. Buchanan TM, et al: Brucellosis in the United States 1960-1972. Med 53:403-439, 1974
3. Young EJ, Suvannaparrat U: Brucellosis outbreak attributed to ingestion of unpasteurized goat cheese. Arch Intern Med 135:240-243, 1975
4. Fox MD, Kaufman AF: Brucellosis in the United States, 1965-1974. J Infect Dis 136:312-316, 1977

Clostridium botulinum

1. Center for Disease Control: Botulism in the United States, 1899-1977. Handbook for Epidemiologists, Clinicians, and Laboratory Workers, CDC, Atlanta, May 1979, pp 12-13

2. Cherington M: Botulism. Ten-year experience. Arch Neurol 30:432-437, 1974
3. Koenig MG, Drutz DJ, Mushlin AI, et al: Type B botulism in man. Am J Med 42:208-219, 1967
4. Koenig MG, Spichard A, Cardella MA, et al: Clinical and laboratory observations of type E botulism in man. Med 43:517-545, 1964
5. Dowell VR Jr, McCroskey LM, Hatheway CL, et al: Coproexamination for botulinum toxin and Clostridium botulinum - A new procedure for laboratory diagnosis of botulism. JAMA 238:1829-1832, 1977

Clostridium perfringens

1. Bryan FL: What the sanitarian should know about Clostridium foodborne illness. J Milk Food Technol 32:381-389, 1969
2. Lowenstein MS: Epidemiology of Clostridium perfringens food poisoning. N Engl J Med 286(19):1026-1027, 1972

Escherichia coli

1. Marier R, Wells JG, Swanson RC, Callahan W, Mehlman IJ: An outbreak of enteropathogenic Escherichia coli foodborne disease traced to imported French cheese. Lancet 2:1376-1378, 1973
2. Sack RB: Human diarrheal disease caused by enterotoxigenic Escherichia coli. Ann Rev Microbiol 29:333-353, 1975

Salmonella

1. Aserkoff B, Schroeder SA, Brachman PS: Salmonellosis in the United States--A five-year review. Am J Epidemiol 92:13-24, 1970
2. Bryan FL: What the sanitarian should know about salmonellae and staphylococci in non-dairy foods. II. Salmonellae. J Milk Food Technol 31:131-140, 1968
3. Cohen ML, Blake PA: Trends in foodborne salmonellosis outbreaks: 1963-1975. J Food Protection 40:798-800, 1977

Shigella

1. Donadio JA, Gangarosa EJ: Foodborne shigellosis. J Infect Dis 119:666-668, 1969
2. Black RE, Craun GF, Blake PA: Epidemiology of common-source outbreaks of shigellosis in the United States, 1961-1975. Am J Epidemiol 108:47-52, 1978

Staphylococcus

1. Bryan FL: What the sanitarian should know about salmonellae and staphylococci in non-dairy foods. I. Staphylococci. J Milk Food Technol 31:110-116, 1968
2. Merson MH: The epidemiology of staphylococcal foodborne disease. Proceedings of the Staphylococci in Foods Conference, Pennsylvania State University, University Park, Pennsylvania, 1973, pp 20-37
3. Minor TE, Marth EH: Staphylococcus aureus and staphylococcal food poisoning. J Milk Food Technol 34:21-39, 77-83, 227-241, 1972, 35:447-476, 1973

Group A Streptococcus

1. Hill HR, Zimmerman RA, Reid GVK, Wilson E, Kitton RM: Foodborne epidemic of streptococcal pharyngitis at the United States Air Force Academy. N Engl J Med 280:917-921, 1969
2. McCormick JB, Kay D, Hayes M, Feldman RA: Epidemic streptococcal sore throat following a community picnic. JAMA 236:1039-1041, 1976

Vibrio cholerae

1. Finkelstein RA: Cholera. CRC Critical Reviews in Microbiology 2(4):553-623, 1973
2. Gangarosa EJ, Mosley WH: Epidemiology and surveillance of cholera: Cholera, edited by Barua D, Burrows W. Philadelphia, London, Toronto, WB Saunders Co., 1974, p 381
3. Center for Disease Control: Follow-up on Vibrio cholerae Infection--Louisiana. Morbidity Mortality Weekly Rep 27(39):367, 1978

Vibrio parahaemolyticus

1. International Symposium on Vibrio parahaemolyticus, September 17-18, 1973, Fujino, Sakaguchi G, Sakazaki R, Takeday (ed). Saikon Publishing Co., Ltd., Tokyo, Japan, 1974
2. Barker WH: Vibrio parahaemolyticus outbreaks in the United States. *Lancet* 1:551-554, 1974
3. Barrow GI, Miller DC: Vibrio parahaemolyticus and seafoods. *Soc Appl Bacteriol Symp Ser* 4:181-195, 1976
4. Dadisman TA Jr, Nelson R, Molenda JR, et al: Vibrio parahaemolyticus gastroenteritis in Maryland. I. Clinical and epidemiologic aspects. *Am J Epidemiol* 96:414-426, 1972

CHEMICAL

Heavy Metal

Cadmium

1. Baker TD, Hafnew WG: Cadmium poisoning from a refrigerator shelf used as an improvised barbecue grill. *Public Health Rep* 76:543-544, 1961

Copper

1. Hopper SH, Adams HS: Copper poisoning from vending machines. *Public Health Rep* 73:910-914, 1958
2. Semple AB, Parry WH, Phillips DE: Acute copper poisoning: an outbreak traced to contaminated water from a corroded geyser. *Lancet* 2: 700-701, 1960

Tin

1. Barker WH, Runte V: Tomato juice-associated gastroenteritis. Washington and Oregon, 1969. *Am J Epidemiol* 96:219-226, 1972

Zinc

1. Brown MA, Thom JV, Orth GL, et al: Food poisoning involving zinc contamination. *Arch Environ Health* 8:657-660, 1964

Ichthyosarcotoxin

1. Hughes JM, Merson MH: Fish and shellfish poisoning. *N Engl J Med* 295:1117-1120, 1976

Ciguatoxin

1. Barkin RM: Ciguatera poisoning. A common-source outbreak. *South Med J* 67(1):13-16, 1974
2. Halstead BW: Poisonous and venomous marine animals of the world. Princeton, The Darwin Press, 1978, pp 325-402

Puffer Fish (tetrodotoxin)

1. Halstead BW: Poisonous and venomous marine animals of the world. Princeton, The Darwin Press, 1978, pp 437-548
2. Torda TA, Sinclair E, Ulyatt DB: Puffer fish (tetrodotoxin) poisoning: Clinical record and suggested management. *Med J Aust* 1:599-602, 1973

Scombrototoxin

1. Halstead BW: Poisonous and venomous marine animals of the world. Princeton, The Darwin Press, 1978, pp 417-435
2. Kimata M: The histamine problem: Fish as Food, edited by Borgstrom E, New York, Academic Press, 1961, pp 329-352
3. Merson MH, Baine WB, Gangarosa EJ, Swanson RC: Scombroid fish poisoning: outbreak traced to commercially canned tuna fish. *JAMA* 228: 1268-1269, 1974

Monosodium Glutamate

1. Schaumburg HH, Byck R, Gerstl R, Mashman JH: Monosodium L-glutamate; its pharmacology and role in the Chinese restaurant syndrome. *Science* 163:826-828, 1969

Mushroom Poison

1. Wieland T, Wieland O: The toxic peptides of Amanita species in Kadis S, Ciegler A, Alj SJ (eds): Microbial toxins, Vol 8 Fungal toxins, New York and London, Academic Press, 1972, pp 249-280
2. Benedict RG: Mushroom toxins other than Amanita, Kadis S, Ciegler A, Alj SJ: Microbial toxins, Vol 8 Fungal toxins, New York and London, Academic Press, 1972, pp 281-320
3. Tyler VE: Poisonous mushrooms: Progress in Chemical Toxicology. Vol 1, edited by Stolman A, New York, Academic Press, 1963, pp 339-384
4. Becker CE, et al: Diagnosis and treatment of Amanita phalloides type mushroom poisoning - use of thiotic acid. West J Med 125:100-109, 1976

Parylytic and Neurotoxic Shellfish Poison

1. Music SI, Howell JT, Brumback CL: Red tide: its public health implications. J Fla Med Assoc 60:27-29, 1973
2. Halstead BW: Poisonous and venomous marine animals of the world. The Darwin Press, Princeton, 1978, pp 43-78
3. Hughes JM, Merson MH: Fish and shellfish poisoning. N Engl J Med 295:1117-1120, 1976

PARASITIC

Anisakidae

1. Chitwood MD: Nematodes of medical significance found in market fish. Am J Trop Med Hyg 19:599-602, 1970

T. spiralis

1. Gould SE: Trichinosis in man and animals. Springfield, Ill, Charles C. Thomas, 1970
2. Zimmerman WJ, Steele JH, Kagan IG: Trichinosis in the U.S. population 1966-1970--prevalence and epidemiologic factors. Health Services Rep 88:606-623, 1973


T. gondii

1. Kean BH, Kimball AC, Christensen WN: An epidemic of acute toxoplasmosis. JAMA 208:1002-1004, 1969

VIRAL

Hepatitis A

1. Cliver DO: Implications of foodborne infectious hepatitis. Public Health Rep 81:159-165, 1966
2. Gravelle CR, Hornbeck CL, Maynard JE, et al: Hepatitis A: Report of a common-source outbreak with recovery of a possible etiologic agent. II. Laboratory studies. J Infect Dis 131:167-171, 1975
3. Leger RT, Boyer KM, Pattison CP, et al: Hepatitis A: Report of a common-source outbreak with recovery of a possible etiologic agent. I. Epidemiologic studies. J Infect Dis 131:163, 1975



STATE EPIDEMIOLOGISTS AND STATE LABORATORY DIRECTORS

The State Epidemiologists are the key to all disease surveillance activities and their contributions to this report are gratefully acknowledged. In addition, valuable contributions are made by State Laboratory Directors.

STATE	STATE EPIDEMIOLOGIST	STATE LABORATORY DIRECTOR
Alabama	Thomas J Chester, MD	James L Holston, Jr, DrPH
Alaska	John P Middaugh, MD	Harry J Colvin, PhD
Arizona	Philip M Hotchkiss, DVM, Acting	Jon M Counts, DrPH
Arkansas	John P Lofgren, MD	Robert T Howell, DrPH
California	James Chin, MD	John M Heslep, PhD
Colorado	Richard S Hopkins, MD	C D McGuire, PhD
Connecticut	Anthony V Sardinias, MPH	Jesse Tucker, PhD
Delaware	Ernest S Tierkel, VMD	Mahadeo P Verma, PhD
District of Columbia	Martin E Levy, MD	James B Thomas, PhD
Florida	Robert A Gunn, MD	Nathan J Schneider, PhD
Georgia	R Keith Sikes, DVM	Frank M Rumph, MD
Guam	Robert L Haddock, DVM	Pedro L G Santos
Hawaii	Ned H Wiebeaga, MD	Albert I Oda
Idaho	Fritz R Dixon, MD	D W Brock, DrPH
Illinois	Byron J Francis, MD	Hugh-Bert Ehrhard, DrPH
Indiana	Charles L Barrett, MD, Acting	Josephine Van Fleet, MD
Iowa	Laverne A Wintermeyer, MD	W J Hausler, Jr, PhD
Kansas	Donald E Wilcox, MD	Roger H Carlson, PhD
Kentucky	Joseph W Skaggs, DVM, Acting	B F Brown, MD
Louisiana	Charles T Caraway, DVM	Henry Bradford, PhD
Maine	William S Nersesian, MD, Acting	Howard E Lind, PhD
Maryland	Ebenezer Israel, MD, Acting	J Mehsen Joseph, PhD
Massachusetts	Nicholas J Fiumara, MD	William M McCormack, MD
Michigan	Norman S Hayner, MD	George R Anderson DVM
Micronesia*	Masao Kumangai, MD	
Minnesota	Andrew G Dean, MD	C Dwayne Morse, DrPH
Mississippi	Durward L Blakey, MD	R H Andrews, MS
Missouri	H Denny Donnell, Jr, MD	Elmer Spurrier, DrPH
Montana	Judy Gedrose, RN, MN, Acting	Douglas Abbott
Nebraska	Paul A Stoesz, MD	John Blosser
Nevada	William M Edwards, MD	Paul Fugazzotto, PhD
New Hampshire	Vladas Kaupas, MD	Robert A Miliner, DrPH
New Jersey	William E Parkin, DVM	Bernard F Taylor, PhD
New Mexico	Sandra C Lapham, MD, Acting	Aaron Bond, MPH
New York State	Richard Rothenberg, MD	David O Carpenter, MD
New York City	Stephen M Friedman, MD	Bernard Davidow, PhD
North Carolina	Martin P Hines, DVM	Mildred A Kerbaugh
North Dakota	Kenneth Mosser	A A Gustafson
Northern Mariana Islands*	Frank T Palacios, MD	
Ohio	Thomas J Halpin, MD	Charles C Croft, ScD
Oklahoma	Mark A Roberts, PhD	Gary McKee, PhD
Oregon	John A Googins, MD	William Murphey, PhD
Pennsylvania	Ernest J Witte, VMD	Vern Pidcoe, DrPH
Puerto Rico	Antonio Hernandez, MD	Jose L Villamil
Rhode Island	Gerald A Faich, MD	Raymond G Lundgren, PhD
South Carolina	Richard L Parker, DVM	Arthur F DiSalvo, MD
South Dakota	James D Corning, BA	A Richard Melton, DrPH
Tennessee	Robert H Hutcheson, Jr, MD	Michael W Kimberly, DrPH
Texas	Charles R Webb, Jr, MD	Charles Sweet, DrPH
Utah	Richard E Johns, Jr, MD	Francis M Urry, PhD
Vermont	Richard L Vogt, MD	Dymitry Pomer, DVM
Virginia	Grayson B Miller, Jr, MD	Frank W Lambert, DrPH
Virgin Islands	John N Lewis, MD	Norbert Mantor, PhD
Washington	Jack Allard, PhD†	Jack Allard, PhD
West Virginia	Loretta E Haddy, MS	John W Brough, DrPH
Wisconsin	Jeffrey P Davis, MD	Ronald Laessig, PhD, Acting
Wyoming	Martin D Skinner, MD	Donald T Lee, PhD

*Formerly Trust Territory of the Pacific Islands

†Dual Assignment

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
PUBLIC HEALTH SERVICE
CENTERS FOR DISEASE CONTROL
ATLANTA, GEORGIA 30333

OFFICIAL BUSINESS

POSTAGE AND FEES PAID
U.S. DEPARTMENT OF HHS
HHS 396

